



The Open Biotechnology Journal

Content list available at: www.benthamopen.com/TOBIOTJ/

DOI: 10.2174/1874070701812010056



REVIEW ARTICLE

Microbial Diversity of *Mer* Operon Genes and Their Potential Roles in Mercury Bioremediation and Resistance

Martha M. Naguib¹, Ahmed O. El-Gendy² and Ahmed S. Khairalla^{2,*}

¹Department of Biotechnology and Life Sciences, Faculty of Post Graduate Studies for Advanced Sciences, Beni-Suef University, Beni-Suef 62511, Egypt

²Department of Microbiology and Immunology, Faculty of Pharmacy, Beni-Suef University, Beni-Suef 62511, Egypt

Received: October 27, 2017

Revised: February 23, 2018

Accepted: March 16, 2018

Abstract:

Background:

Mercury is a toxic metal that is present in small amounts in the environment, but its level is rising steadily, due to different human activities, such as industrialization. It can reach humans through the food chain, amalgam fillings, and other sources, causing different neurological disorders, memory loss, vision impairment, and may even lead to death; making its detoxification an urgent task.

Methods:

Various physical and chemical mercury remediation techniques are available, which generally aim at: (i) reducing its mobility or solubility; (ii) causing its vaporization or condensation; (iii) its separation from contaminated soils. Biological remediation techniques, commonly known as bioremediation, are also another possible alternative, which is considered as cheaper than the conventional means and can be accomplished using either (i) organisms harboring the *mer* operon genes (*merB*, *merA*, *merR*, *merP*, *merT*, *merD*, *merF*, *merC*, *merE*, *merH* and *merG*), or (ii) plants expressing metal-binding proteins. Recently, different *mer* determinants have been genetically engineered into several organisms, including bacteria and plants, to aid in detoxification of both ionic and organic forms of mercury.

Results:

Bacteria that are resistant to mercury compounds have at least a mercuric reductase enzyme (MerA) that reduces Hg⁺² to volatile Hg, a membrane-bound protein (MerT) for Hg⁺² uptake and an additional enzyme, MerB, that degrades organomercurials by protonolysis. Presence of both *merA* and *merB* genes confer broad-spectrum mercury resistance. However, *merA* alone confers narrow spectrum inorganic mercury resistance.

Conclusion:

To conclude, this review discusses the importance of mercury-resistance genes in mercury bioremediation. Functional analysis of *mer* operon genes and the recent advances in genetic engineering techniques could provide the most environmental friendly, safe, effective and fantastic solution to overcome mercuric toxicity.

Keywords: Mercury, Mercury toxicity, Biogeochemical cycle, Mercury remediation, Resistance mechanisms, *Mer operon*.

1. INTRODUCTION

For a long time, the term “heavy metals” had been widely used for metals associated with contamination and eco

* Address correspondence to this author at the Department of Microbiology and immunology, Faculty of Pharmacy, Beni-Suef University, Beni-Suef 62511, Egypt, Tel: +2-012-234-76015; E-mails: ahmed.elgendy@pharm.bsu.edu.eg, ahmedkhairalla@pharm.bsu.edu.eg

-toxicity. The International Union of Pure and Applied Chemistry (IUPAC) recommended using the term “toxic metal” as an alternative to “heavy metal” [1]. Toxic metals are stable and persistent environmental contaminants [2]. Many metals such as mercury, cadmium, chromium, zinc, lead, copper, arsenic *etc.*, used in different industries, are releasing its toxic ions and introducing it into the ecosystem leading to toxic effects, affecting humans, animals, plants, and microbial communities [3]. Also, some toxic metals naturally exist in very low concentrations in the ecosystem and are required in trace amounts as nutrients by microbial communities but in relatively higher concentration, they form toxic complexes on the biological cell [3, 4].

Mercury is the 16th rarest element on the earth [5] and considered as one of the mobile and toxic metals that exist naturally in low concentrations in the environment, and can be changed between different forms. It is the only metal to be liquid at room temperature. It can also exist as gas due to its high vapor pressure [6 - 8].

It is a major environmental pollutant especially methylmercury (MeHg) form in the aquatic regions. It can accumulate in biota so can reach and affect both wildlife and human seriously. It is one of the environmentally stable and persistent toxins for long periods, also it can be accumulated in different biological tissues [9, 10].

This review aims to cover all aspects related to the environmental biogeochemical cycle of mercury, the roles of *mer* genes in microbial adaptation to mercury, and potential bacterial remediation strategies of this toxic metal.

2. MERCURY BIOGEOCHEMICAL CYCLE

Environmental mercury cycle, as illustrated in Fig. (1), is usually facilitated biotically and abiotically between soils, water and atmosphere [11]. Mercury exists in the atmosphere in gaseous, particulates, and aqueous soluble forms [12] but gaseous form represents about 95% of atmospheric mercury [13] and it remains in the atmosphere for long periods. So, it can reach the far distance that should be considered as a huge environmental concern [12, 14, 15].

2.1. Oxidation Processes

In the atmosphere, Hg abiotically oxidized to Hg^{+2} through photo-oxidation reactions, mediated by O_2 through its interaction with hydrogen peroxide, ozone, sulfhydryl compounds, free radicals as Br, and by UV-B in presence of Cl_2 and photoreactive compounds as benzoquinone in presence of water droplets [11, 12, 16].




Fig. (1). Environmental mercury biogeochemical cycle [10].

Biotic Hg oxidation (bio-oxidation) in aerobic and phototrophic microorganisms is catalyzed by hydroperoxidases, *katG* and *katE* [13] or other oxidases [17]. Algae, plants and animals catalase and peroxidases are also capable of oxidizing Hg [17 - 19].

Anaerobic Hg oxidation mechanism by *Desulfovibrio desulfuricans* ND132 bacteria is unknown as obligate

anaerobes do not carry such catalase or peroxidase genes but Colombo, Ha *et al.* 2013 suggested an alternative oxidation pathway influenced by reactive functional thiol groups of different anaerobic bacteria [17, 19] consistent with previous studies in which thiol moiety of organic compounds such as glutathione [20] and reduced humic acid [21] can bind and oxidize Hg.

2.2. Reduction Processes

Biotic Hg^{+2} reduction to Hg occurs by bacterial mercuric reductase enzyme encoded by *merA* gene on *mer* operon (described later) [8, 11, 22], anaerobic *Geobacter sulfurreducens* PCA, *Geothrix fermentans* H5, *Cupriavidus metallidurans* AE104, *Shenwella oneidensis* MR-1 and *Geobacter metallireducens* GS-15 was found to reduce Hg^{+2} to Hg independent on the *mer* system but its need an electron acceptors and electron donors to suggest activity of respiratory electron transport chains. Its activity is effective at too low Hg^{+2} concentrations compared to amounts required for *mer* operon induction. Anaerobic bacteria showed dual role in the Hg redox cycle by both oxidizing Hg and reducing Hg^{+2} by unidentified reduction system other than *mer* system [19, 22].

The abiotic Hg^{+2} reduction is done not only by photochemical reactions but also *via* dark reactions [23] using organic matter free radicals as fulvic [24] and humic acid-associated free radicals [18].

2.3. Methylation Process

Biotic Hg^{+2} methylation is a natural bacterial process mainly occurred in seawater and coastal environment sediments [25], invertebrate digestive tracts, thawing permafrost soils, and extreme environments [26] through anaerobic methylators as Sulfate-Reducing Bacteria (SRB) [27 - 30], Iron Reducing Bacteria (IRB) and methanogens [27, 31, 32]. Two-gene cluster, *hgcA* and *hgcB* encode a putative corrinoid protein facilitating methyl transfer and a ferredoxin carrying out corrinoid reduction, were reported to be involved in mercury methylation process [27]. According to Podar *et al.* 2015, *hgcA* and *hgcB* genes were found in nearly all anaerobic environments but not in aerobic and not in human and mammalian microbiomes, reducing the expected risk of Hg methylation by microbiomes [26].

Another unrecognized oxidation/methylation pathway in the mercury cycle by anaerobic bacteria as in *D. desulfuricans* ND132 and *G. sulphurreducens* PCA [17, 33] produces MeHg using dissolved Hg as their sole Hg source. Further investigation is required for detecting the reactions involved in and the connection between Hg methylation, oxidation and produced toxic MeHg exportation out of cells as *D. desulfuricans* lacks the *mer* operon system that detoxifies organic/inorganic mercury [17].

Moreover, abiotic Hg^{+2} methylation carried out chemically [34] with the help of humic and fulvic acids, carboxylic acids, and compounds as fungicides or antifouling agents [11, 35]. It was found that sunlight played a minor and slower role in methylation reaction [36].

2.4. Demethylation Process

Biotic demethylation occurs simultaneously in the methylation sites as a reverse process [34]. It is catalyzed through reduction [37] or oxidation processes [38]. Aerobic reductive demethylation of CH_3Hg^+ occurs through *mer* operon functions (*merA* and *merB*) forming CH_4 and Hg [37]. Formed Hg will be evaporated into the air, and the cycle is repeated [39, 40]. Anaerobic reductive demethylation, *Geobacter bemidjiensis* Bem is an iron-reducing bacterium capable of simultaneously both methylating Hg^{+2} and degrading MeHg, due to the the presence of homologues of an organomercurial lyase and a mercuric reductase [41]. Oxidative demethylation occurs in more anaerobic conditions yielding CO_2 and small amounts of Hg^{+2} with an unknown mechanism [38, 39]. Produced Hg^{+2} may be available for re-methylation process or reduction to its vapor form [11]. Both anaerobic methylation and demethylation processes are affected by environmental dissolved organic matter, iron-sulfate biogeochemistry, and Hg^{+2} concentration [17, 27, 42].

Abiotic demethylation of MeHg affected by photo-degradation, especially UV-A and UV-B, at a wavelength range of 200-400 nm [11, 43, 44]. A study on Hg cycling found that dissolved MeHg concentration was decreased in daylight and increased in non-daylight suggesting that photo-degradation in water has a major role in methylation/demethylation processes aquatic systems [45].

All mercury forms (Hg , Hg^{+2} , and CH_3Hg^+) are interconvertible and can be introduced into the aquatic system [8]. Their concentration in the aquatic system depends mainly on reduction, methylation, and demethylation ratio which

depend on the microbial community [6].

CH_3Hg^+ had a different pathway as bio-accumulator in the food chain. Its concentration is higher in top food chain organisms due to CH_3Hg^+ biomagnifications [46]. As CH_3Hg^+ concentration is higher in predatory animals such as beluga and polar bears than marine mammals which have higher concentrations of Hg than freshwater fish and overland mammals, due to their higher position in marine food webs [47]. Marine and freshwater fishes are still the main sources of dietary Hg exposure for humans [6, 47]. CH_3Hg^+ is the main specie absorbed in the gut; it enters the bloodstream and distributes to body tissues and organs [47]. In fish, CH_3Hg^+ first accumulates in the viscera (kidney, spleen, and liver) and later redistributed to other tissues as muscles and brain tissues. Hence, mammals and birds can demethylate CH_3Hg^+ . So, a large amount of accumulated Hg in liver and kidney are found in an inorganic form [47].

3. MERCURY BIOREMEDIATION

It is a remediation technique using a wide range of living organisms (algae, fungi, yeasts, plants and bacteria) or their enzymes. Bioremediation is preferred as environmental friendly, promising technique. Microbial remediation using bacteria is widely used as they can be easily cultivated, grow faster and can accumulate metals in different conditions [48 - 51]. Moreover, different Gram-negative and positive bacterial isolates can resist, accumulate, adsorb and transform toxic mercury forms to less toxic forms by different mechanisms. These bacteria are named Mercury Resistant Bacteria (MRB) [8, 12, 48]. Bioremediation using bacterial strains showed promising results, reached 76.4% compared to other remediation techniques in removing mercury pollutant leached from spent fluorescent lamps [52].

3.1. Phytoremediation

Green plants or its associated microorganisms are used to remove or destroy contaminants from the soil. The insertion of bacterial mercury resistance genes as (*merP*, *merC*, *merF* and *merT*) encodes for different transporter proteins, (*merA*) encodes for mercury reductase or (*merB*) encodes for organomercurial lyase into plant cells after their sequences modification according to preferred plant codons. Genes inserted as *merB* remove organic mercury by protonolysis of C-Hg to Hg^{+2} while *merA* helps in the reduction of Hg^{+2} by the formation of volatile elemental mercury Hg which is then volatilized out of plant cells [53 - 57]. Different plant types and species were engineered as *Arabidopsis thaliana* as described in Table 1, yellow poplar [58], tobacco [59, 60], peanut [61], salt marsh cordgrass [62], rice [63] and eastern cottonwood [64, 65] that showed successful high resistance levels than their wild types.

Hence, genetically engineered plants can get rid of ionic and organic mercury by phytovolatilization. However, phytovolatilization major concern is the release of mercury vapors back to the environment, but this can be reduced by increasing efficacy of phytoextraction/phytosequestration. This could be achieved by transforming plants with certain bacterial genes as *merP*, *merC*, *merF* and *merT* [53]. For example, the increase of Hg^{+2} bioaccumulation in transgenic tobacco by expressions of the *merP* gene producing bacterial Polyphosphate Kinase (PPK) [54, 66].

An MRB Enterobacter strain exhibited a novel property of Hg immobilization by synthesis of nanoparticle Hg. The strain could intracellularly synthesize Hg nano-particles sized 2–5-nm [67].

Table 1. Effect of different *mer* genes, inserted and expressed in different regions of *A. thaliana* plant, on mercury phytoremediation.

Expression	Inserted gene	Effect on Resistance	References
Cytoplasm	<i>merB</i>	Increase resistance of phenylmercuric acetate PMA to 2 μM	[157].
Cytoplasm	<i>merA</i>	Increase resistance of Hg^{+2} to 100 Mm	[56].
Cytoplasm	<i>merA/B</i>	Increase resistance of organic and inorganic Hg to 5–10 μM	[158].
Cell membrane	<i>merP</i> (<i>ppk</i>)	Increase resistance of Hg^{+2} to 10 μM	[159].
Cell membrane	<i>merC</i>	Increase resistance of Hg^{+2} to 10 μM	[160].
Cytoplasm	<i>merE</i>	Increase of inorganic and organic-Hg accumulation	[55].

4. MERCURY RESISTANCE MECHANISMS

To adapt to toxic metals in the environment, bacteria and other organisms have developed different resistance mechanisms as a defense systems against these toxic materials. This defense systems that help bacteria to eliminate toxic materials from their growth medium includes:

1. Mercury bioaccumulation whether by simultaneous synthesis of mercury as nanoparticles [67] or by Hg^{+2} binding to carboxyl phosphates, hydroxyl, thio, or pyridine functional groups located on some bacterial cell wall [49] so, mercury is trapped and can't be vaporized back into the environment [67],
2. Sequestration and chelation of mercury using intracellular binding Metallothionein protein [68], a cysteine-rich protein able to bind mercury ions to form Mercury-cysteine complex or extracellular polysaccharides in the cell wall [69, 70] as mercury compounds have high ability to bind with thiols of bacterial cysteine and reduce mercury toxicities [11],
3. Blocking mercury entry into cells through permeability barriers [11, 71],
4. Efflux and volatilization to convert toxic ionic mercury, Hg^{+2} , to much less toxic, elemental mercury, Hg through genetic manner [70], reductase enzymes as those expressed by the help of mercury resistance (*mer*) operon that will be described later [3, 70, 72] or through cytochrome c oxidase enzymes [73].

For these valuable mechanisms, different bacterial strains and other biological systems were engineered to be used for remediation and monitoring of environmental hazards such as increasing the bioaccumulation of Hg^{+2} by expression of the bacterial Polyphosphate Kinase (PPK) in transgenic tobacco [66].

5. COMPONENTS AND FUNCTIONS OF *MER* OPERON IN MERCURY RESISTANT BACTERIA (MRB)

All *mer* determinants (*mer* operon) are widely distributed by both Horizontal (HGT) and Vertical Gene Transfer (VGT) which explain their presence in different bacterial populations [70]. It was found located on chromosomal DNA [74], mobile elements as plasmids [40, 75], transposons as components of the Tn21 [22, 76] and Tn501 [46], or on integrons [77]. These *mer* determinants were identified in a wide range of previously isolated gram-negative [78] as seen in Fig. (2) and Gram-positive [79] bacterial strains from clinical [80, 81] and environmental [82, 83] samples. Genes in *mer* operon express different enzymes that can transform toxic to less toxic mercury forms as organomercuriallyase and mercuric reductase as illustrated in Table 2. In addition to mercury detoxification, some MRB can also detoxify other metals [84].

These *mer* determinants are classified into two types; narrow-spectrum that detoxifies only inorganic mercury through the main *merA* gene or broad-spectrum that detoxifies both organic and inorganic mercury through *merA* and *merB* genes [8, 85, 86]. The *mer* operon is composed of the operator, promoter, regulator genes, and functional genes such as *merR*, *merP*, *merT*, *merD*, *merA*, *merF*, *merC*, *merE*, *merH*, *merG* and *merB*. All these genes code for different proteins that participate in the detection, transportation and reduction or methylation of mercury ions [7, 75, 87, 88].

Table 2. All *mer* operon genes and their expressed proteins.

Gene	Gene Location in Operon	Protein Encoded	Protein Location	Function	References
<i>merA</i>	Immediately after transport genes	Mercuric reductase	Cytoplasm	Reduction of Hg^{+2} to Hg	[8, 22].
<i>merB</i>	Immediately after transport genes	Organomercuriallyase	Cytoplasm	Lysis of C-Hg bond	[8, 99].
<i>merF</i>	May be in plasmid between <i>merP</i> and <i>merA</i>	Transporter protein	Inner membrane	Narrow and broad spectrum system transporters of mercuric.	[90, 92].
<i>merE</i>	Downstream of <i>merA</i> and <i>merB</i>	Transporter protein	Inner protein	Uptake of inorganic and organo-mercurials (MeHg) into cytoplasm	[55, 89, 69].
<i>merG</i>	Between <i>merA</i> and <i>merB</i>	Organomercurial compounds resistance protein	Periplasm	Resistance to phenyl-mercury by efflux mechanism (reduce their cellular permeability)	[11, 89, 86]
<i>merP</i>	Upstream <i>merA</i>	Periplasmic Hg^{+2} binding protein	Periplasm	Transfer of Hg^{+2}	[8, 11, 76].
<i>merT</i>	Upstream <i>merA</i>	Transporter protein	Inner membrane	Transporters of mercury in both narrow (Hg^{+2}) and broad (phenyl mercury) spectrum systems.	[89, 90].
<i>merD</i>	Downstream <i>merA</i>	Regulator protein	Cytoplasm	Negatively regulates the <i>mer</i> operon	[109, 110,].
<i>merC</i>	Upstream <i>merA</i>	Transporter protein	Inner membrane	Narrow and broad spectrum system transporters of mercuric.	[90],
<i>merR</i>	Before transport genes	Regulator protein	Cytoplasm	Positively regulates the <i>mer</i> operon	[11, 22, 76, 89, 106].

Gene	Gene Location in Operon	Protein Encoded	Protein Location	Function	References
<i>merH</i>	Immediately adjacent to <i>merA</i>	Mercury trafficking protein (metallochaperone)	Cytoplasm	Trafficking of mercury	[95, 161].
<i>merI</i>	Located 3' to <i>merA</i>	Unknown		Have no role in mercury resistance or regulation	[95, 161].

5.1. Hg⁺² Binding and Transportation Genes

merT, *merP*, *merC*, *merE*, *merF*, and newly discovered *merH* express different proteins that have different roles in mercury transportation as shown in Fig. (2) [89].

merT, expresses an inner membrane (cytoplasmic) proteins [22], helps in the uptake of organic phenylmercury [90] and inorganic mercury transport into the cytoplasm [91]. MerT protein has three transmembrane regions with cysteine pair located in its first transmembrane region Cys-Cys [90].

merP, expresses a periplasmic protein [22] that has two cysteine residues to help in replacing the nucleophiles (Cl⁻) linked to Hg⁺² so, Hg⁺² could bind to merP cysteine residues then transferred to other two cysteine residues on merT protein located on cytoplasmic membrane [8, 11, 76] then passed to other pair of merT's cytosolic cysteines residues. Hg⁺² enters the cytoplasmic membrane where cytosolic thiols (cysteines or glutathions) compete with merT's cytosolic cysteines to bind with Hg⁺² to be ready for merA activity [11].

merT and *merP* deletion from transposon Tn501 lead to Hg⁺² sensitivity while the expression of Tn501 *merT* and *merP* in the absence of mercuric reductase causes Hg⁺² supersensitivity [22]. Moreover, mutations in both *merT* and *merP* increase Hg⁺² concentration required for induction of merA-lacZ transcriptional fusions. So, *merP* is important for Hg⁺² resistance [22]. However, Sone and Nakamura *et al.* showed that absence of *merP* in presence of other alternative transporters *merC*, *merE*, *merF* and *merT* would increase both inorganic and organic mercury transportation. While the presence of *merP* with all other transporters did not cause any difference in organic mercury transportation but increased the inorganic transportation [90]. MerE, MerT, MerC and MerF are broad-spectrum mercury transporter proteins that can transport organic phenylmercury and inorganic Hg⁺² into cells [11, 90 - 92].

MerC, expresses an inner membrane (cytoplasmic) protein [22], that is involved in both inorganic and organic phenylmercury [90] uptake across the cytoplasmic membrane till it reaches active site reductase [70]. *MerC* protein was found to have roles in mercury accumulation in *Arabidopsis thaliana* [93].

Absence or mutation of *merC* could show no effects on mercury transportation and resistance in case of presence of *merT* and *merP* as shown in Tn501 and Tn5053. However, in some bacterial strains when both *merT* and *merP* do not exist, *merC* can act alone as the main Hg⁺² transporter [22, 76]. Other new findings showed that *merC* is more preferred in Hg⁺² transportation than *merE*, *merF* and *merT*. Moreover, *merC* was more efficient for designing successful mercurial bioremediation system [90].

merF helps in both organic phenylmercury [90] and inorganic Hg⁺² transport across the cytoplasm [11, 92].

merH expresses a membrane protein and according to Schué, it was able to transport Hg⁺² across the inner membrane using its two cysteine residue. Replacing *merT* gene in an *E-coli* strain by *merH* resulted in Hg⁺² MIC reduction, although the strain is still resistant when compared with a control strain which has no transporter proteins [94]. As described by Schelert, a metallochaperone, homolog to C-terminal domain, called TRASH has a role in metal sensing and trafficking that explains *merH* role in trafficking of Hg⁺² to the MerR transcription factor. By non-sense or in-frame gene mutation of *merH*, the new mutant strain was highly sensitive to Hg⁺² and analysis of *merH* deleted operon by mass spectroscopic. They found increased retention of Hg⁺² intracellular that explains the low rate of *mer* operon induction and the need for *merH* in metal trafficking [95].

merE is involved in the transport system of inorganic [89] and organic phenylmercury compounds selectively across the membrane [96]. Organomercurial compounds are lipid-soluble and can pass through the cell membrane by simple diffusion [11, 97] or are transported inside by *merE* or *merG* [98]. Their transport system is poorly understood [96, 97].




Fig. (2). Model of bacterial *mer* operon determinants and expressed genes. Organic methylmercury can enter inside cell through merP, T, and E transporter while, phenylmercury transported by merG protein also both may passively diffused inside the cell. Inorganic mercury transported by merP and T to be reduced by merA(10).

5.2. *mer* Detoxification Genes

merA is the main gene in *mer* operon encodes for mercuric ion reductase enzyme, which is a flavoprotein catalyzes reduction of Hg^{+2} into volatile Hg by using NADPH located in the cytoplasm as source of electrons for the reduction of Flavin Adenine Dinucleotide (FAD) [76, 99, 100]. Hg is released out of the bacteria through cell membrane due to its lipid solubility without efflux system [11, 22, 75, 76, 101]. The released Hg back to the environment causes repetition of mercury cycle. To overcome this problem, a new study used an engineered bacteria to express polyphosphate, a chelator for divalent metal, in addition to *mer* operon determinants to overcome metal volatilization [102], help reducing environmental re-pollution. *merA* has an amino-terminal domain (NmerA) Fig. (3), which is homologous to *merP* and contains a pair of cysteines that directly remove Hg^{+2} from transport proteins cytosolic cysteines to be ready for reduction [11]. Mutations by insertion or deletion in *merA* cause Hg^{+2} hypersensitive strains [22].

merB, codes for organomercurial lyase catalyzes demethylation of organic mercury compounds (alkyl and aryl) via cleavage of its C-Hg bond releasing a protonated organic moiety (as methane (CH_4) in case of MeHg) and Hg^{+2} which is then reduced to Hg by merA using same NADPH-dependent mechanism [11, 78, 99, 103, 104].

5.3. Mechanism of Organomercurials Protonolysis by Merb

Organomercuriallyase acts as a monomer containing four cysteine residues Cys117, Cys 96, Cys 159, and Cys160. Asp99, Cys96, and Cys159 residues of the enzyme and water molecule all are involved in protonolysis reaction [11, 103]. MeHg forms bond with Cys96, while the Cys159 site is fully reduced then the organic moiety released through two mechanisms, Fig. (4). In mechanism 1; proton from Cys159 transferred to methyl carbon and Cys159 forms covalent bond with Hg^{+2} . In mechanism 2, Cys159 first transfer proton to Asp99 then forms a covalent bond with Hg^{+2} in presence of methyl group. Asp99 then transfer a proton to the methyl group and the protonated methyl moiety is released. Hg^{+2} is attached to the enzyme by two sulfurs of Cys96 and Cys159, and oxygen from a water molecule [103]. Organomercuriallyase remains attached to Hg^{+2} until two solvent thiols separately bind two Hg^{+2} and removes it [11].

merG helps in cellular permeability of organomercurial compounds (Phenylmercury) through effluxing [11, 96, 98, 105]. Mutation by *merG* deletion, phenylmercury resistance was affected however, Hg^{+2} volatilization activity wasn't affected. So, *merG* have no role in inorganic mercury compounds detoxification [96].




Fig. (3). Illustration for the role of NmerA in Hg^{+2} removal from transport proteins cytosolic cysteine to be reduced by mercuric reductase into Hg (11).

5.4. mer Operon Regulatory Genes

Regulatory genes are *merR* and *merD* [89]. *merR*, a regulatory gene encodes a metalloregulatory protein. It is Hg^{+2} dependent transcriptional repressor-activator that can sense metal concentration and control the expression of other functional *mer* operon genes [11, 106, 107]. It binds with the promoter-operator region (*merOP*) and positively and negatively regulate *mer* operon genes expression in presence or absence of Hg^{+2} , respectively [8, 22, 75, 76, 87, 106]. Hg -free-*merR* (apo-*merR*) undergoes conformational structure changes upon Hg^{+2} binding and converted from repression to activation state Hg -*merR* [108].

merD, a gene encodes for a secondary regulatory protein, present downstream *merA* [109]. MerD protein is expressed in very small amounts and downregulates the *mer* operon [110]. Its absence causes an increase in *mer* operon expression. It binds with the same operator-promoter region (*merOP*) as MerR [8, 22, 76]. It acts as activation antagonist for MerR function. *In vitro* experimental showed that, after Hg^{+2} volatilization by *merA*, the MerR does not give up its bound Hg^{+2} quickly causing *merA* to be active. *merA* expression should be stopped when Hg^{+2} is exhausted because *merA* has an oxidase activity and produces toxic hydrogen peroxide in the absence of Hg^{+2} [11].

6. MER OPERON EVOLUTION, MOBILITY AND DIVERSITY

mer operon is an ancient system [97] may be located on the chromosome [74] and transferred vertically from parents to offspring or on Mobile Genetic Elements (MGE). DNA parts encode proteins that help in its mobility within bacterial genomes or between bacterial cells, facilitating Horizontal Gene Transfer (HGT) independent on reproduction [46, 77, 106, 111 - 114] so, this flexibility in genome [115] can help bacterial adaptation, social interactions and evolution by transferring good genes for the host as the *mer* genes [76, 114, 116 - 119]. MGE can help in the rapid spread of rare, spontaneous resistance mutants to a new bacterial population [114]. As plasmid-borne resistance genes can be originated as point mutations in sensitive bacteria and then transferred when they are flanked by short transposons, picked up by Tn3 family transposons or as mobile cassettes by integrons [76, 114, 120]. *mer* operon as part of different types of group II transposons [121, 122] as Tn2, Tn501, pKLH2, pMERPH and Tn5053 in Gram-negative bacteria [11, 97, 121] and Tn5085, Tn5083 [122, 123], and newly identified Tn6294 [123] in Gram-positive bacteria, All of them encode genes for transposition (*tnpA*) and resolution functions (*tnpR*) [11, 121 - 124].

Successive exposure to mercury for a long time caused *mer* operon persistence and transferred between microbes through HGT and to be evolved rapidly and became more complex as *merA* and species evolved. Evolution happened to adapt with environment, increasing mercury toxicity due to increased industrialization [89] that can explain the

global distribution of *mer* determinants and their associated mobile elements between different bacterial strains [11, 97]. So, *merA* was considered as a biomarker for measuring diversity in Hg detoxification and *mer* operon evolution [89, 125].

A recent study on different *mer* operons of different *Bacillus* species (as tndMER3, tn6294, and others) isolated from thirteen different countries is a good example for *mer* diversity, mobility, evolution and horizontal distribution between different *Bacillus* species worldwide [123].

Genes that express transporter proteins *merP* and *merT* were more common to occur in earliest evolved operons which were less complex while, other alternative transporter genes as *merC*, *merF*, *merE*, and *merH* commonly occurred in more complex operons evolved recently. In organomercurial detoxification system, *merG* gene occurred in more recently evolved operons [89]. Proteins as *merE*, *merC*, *merP*, *merD*, and *merT* were also found to be more likely encoded on plasmids than others so, prefer to be transferred horizontally [89].

A recent research depended on sequencing of four different pQBR mercury resistance plasmids showed that mercury resistance transposons Tn5042 were totally similar in the four plasmids except for one base pair of *merR* in pQBR103 and pQBR44 differed from pQBR55 and pQBR57, indicating that Tn5042 had been transferred between different pQBR plasmids by recombination. They also found that occurrence of pQBR55, pQBR57 and pQBR103 separately in *Pseudomonas fluorescens* SBW25 host resulted in different response to some environmental factors as Hg(II) concentration, although similarity in their *mer* operon suggested the effect of other plasmid-encoded genes [8, 126].




Fig. (4). Mechanisms of MerB catalysis for the Hg-C protonolysis reaction(103). Mechanism 1; Cys159 protonates the methyl group then reacts with Hg⁺². Mechanism 2; hydrogen was attached to the sulfur from Cys159 transferred to Asp99 then Asp99 utilized for methyl group protonation.

6.1. Diversity of *mer* Operon

Several variations in structure and organization of *mer* operons genes are known between both Gram-negative and Gram-positive bacterial strains as in Figs. (5 and 6), for examples:




Fig. (5). Diversity mer operons in Gram-positive bacteria. Arrows indicate the gene product translation direction.

1. *merB* is more common in Gram-positive *mer* operons [123, 127, 128], while *merD* and *merC* or *merF* are more common in Gram-negative *mer* operons [11, 76, 97, 129, 130].
2. Narrow-spectrum *mer* operon genes are highly divergent compared to broad-spectrum operon genes. According to Narita, *et al.* in certain broad-spectrum mercury resistant *Bacillus species*, Polymerase Chain Reaction (PCR) product sizes of *mer* operon genes are identical to that of *Bacillus megaterium* MB1. However, in narrow-spectrum *mer* operon of certain *Bacillus species*, PCR product sizes of the targeted *merP* and *merA* regions are smaller than *merP* and *merA* of the *B. megaterium* MB1 [131].
3. Three different *merB* genes were identified in different *Bacillus species* [127] as in TnMER11-like transposons [123] resistant to different organomercurial compounds and also have multiple *merR*.
4. Transcription direction of *merR* is same as other *mer* operon functional genes in Gram-positive *bacteria mer* operons while, *merR* transcription is divergent from the structural genes in the high-GC Gram-positive *mer* operons and Gram-negative operons except *Pseudoalteromonas haloplanktis* [11, 129, 132, 133].
5. Transposons Tn5084 and Tn5085 are identical in the genetic orientation and contain *merB3*, *merR*, *merE*, *merT*, *merP* and *merA* while, Tn5083 lacks *merR2*, *merB2* and *merB1*, compared to Tn5084 and Tn5085 [122, 123, 132].
6. Newly identified transposon Tn6294 has one *merB*. It is the first transposon to carry *merA* with only one N-terminal mercury binding domain, unlike all other reported broad spectrum *Bacillus species* that carry duplicate N-domain [123].
7. Tn^dMER3, a newly identified deleted transposon, was similar to the fragment in *Bacillus species* that carry *merRETPA* with >90% identity to Tn6294 but with no *merB* and transposonase genes [123].
8. *mer* operon of *Shewanella putrefaciens* pMERPH does not have both *merD* or *merR*, compared to other Gram-negative bacteria suggesting that *merR* may be located elsewhere on the plasmid genes [130].
9. Gene position differs from one operon to another as *merB* present between *merA* and *merD* in pDU1358 [133], while in broad spectrum part of *Pseudomonas stutzeri* pPB located between *merR* and *merT* [134].
10. In *Pseudomonas sp.* Tn502 and Tn512 *mer* operon are related to Tn5053 with exception of *merC* and *urf2M* in newly recognized Tn502 compared to *merF* in Tn512 and Tn5053 [129, 135].

7. CO-RESISTANCE OF MERCURY AND ANTIBIOTICS

mer operons are often part of group II transposons that carry integrons with multiple antibiotic resistance genes [11].

Antibiotics and metals resistance genes are located on the same plasmid [81, 136]. So, mercury exposure can also promote Horizontal Gene Transfer (HGT) of both mercury and antibiotic resistance [91, 125, 137, 138] that explain the increasing challenge in infectious bacteria treatment due to increasing of co-resistance.




Fig. (6). Diversity of *mer* operons in Gram-negative bacteria. Arrows indicate the gene product translation direction.

Six adult monkeys were examined for both mercury and antibiotics resistances of different oral and intestinal bacterial strains before and during the installation, and after the replacement of the amalgam fillings. There was an increase in MRB during the 5 weeks after installation and during the 5 weeks after replacement. MRB was also resistant to one or more antibiotics as streptomycin, kanamycin, tetracycline, ampicillin, and chloramphenicol [81]. In a different study groups exposed to amalgam, MRB isolated from their fecal samples was found to be more resistant to antibiotics than MRB of those never exposed to amalgam [77], suggesting that both antibiotic and mercury resistance genes may be genetically linked [81] and contained within the same genetic mobile element [80] specially, the *Tn21* family of transposons in which the *mer* locus is linked to an antibiotic multi-resistance element [77].

A larger scale study by Pal *et al.*, for the co-occurrence of different metals and antibiotic resistance genes of fully sequenced 2522 bacterial genomes and 4582 plasmids as illustrated in Fig. (7), showed that although metal-antibiotic genes co-resistance was found to be rare on plasmid but, the only metal resistance genes commonly co-occurred with antibiotic-resistant genes on plasmids are mercury resistance genes as seen in Fig. (8). Moreover, about 86% of bacterial genomes contain different metals resistance genes of which 17% co-resistant with antibiotic resistance. Plasmids and genomes with different metals resistance genes were with high probability of carrying antibiotic resistant genes compared to those without metals resistance genes [138].

8. MRB DETECTION METHODS

8.1. Genotypic (Molecular) Techniques

8.1.1. Polymerase Chain Reaction

Simple PCR Specific DNA sequences of the *mer* system determinants could be amplified from the genomic DNA of all the isolates from environmental sources using *mer* determinants designed primers then product can be visualized

under UV on electrophoresis agarose gel as bands by staining with ethidium bromide [101, 139 - 141].

Real-time Reverse Transcriptase-PCR is used to detect not only the presence but also the mRNA expression of the gene [87, 142].




Fig. (7). Overview of the resistance information from (a) 2522 completely sequenced bacterial genomes and (b) 1926 plasmids harboured by those genomes from different environments. (BMRGs) biocide and metal resistance genes. (ARGs) antibiotic resistant genes [138].




Fig. (8). Co-occurrence network of resistance genes on plasmids [138]. The thickness of each connection between two resistance genes is proportional to their co-occurrence times on the same plasmids. ARGs: Antibiotic Resistance Genes; MRGs: Metal Resistance Genes; BRGs: Biocide Resistance Genes; MGEs: Mobile Genetic Elements.

8.1.2. DNA Hybridization Techniques

mer determinants encoding resistance to Hg^{2+} as *merA* or other determinants are used as labeled probes in different hybridization procedures for detecting the presence of *mer* determinants that are complementary to probe nucleotide sequence in mercury resistant bacteria isolated from polluted environments through positive hybridization reaction [139 - 141]. Different hybridization techniques can be applied as Southern [139] and Northern blot hybridization [75], Colony blot hybridization [140, 143], Fluorescent *In Situ* Hybridization (FISH) [144], and microarray that could be used for detection of different multiple gene at once [145] also Restriction Fragment Length Polymorphism (RFLP) could be used [139, 146].

8.1.3. DNA Sequencing

PCR and/or hybridization products can be sequenced to detect its homology to *mer* determinants [139].

8.2. Phenotypic Techniques

8.2.1. Broth and Agar Methods (Direct Bacterial Growth)

Processed environmental samples are added into culture media as Luria-Bertani or nutrient agar (broth) supplemented with 10ppm or greater HgCl_2 to detect visible growth (turbidity or colonies) in media with 10ppm HgCl_2 or more. Grown bacteria are considered as MRB [104, 147, 148]. Optical Density (O.D.) in different time ranges at 600 nm can be measured to detect the growth pattern of tested microorganism in presence of mercury at constant conditions [80, 149].

8.2.2. Hg Utilization (Removal/Reduction) Rate

As only mercury resistant strains can utilize Hg and volatilize it, so the rate of Hg loss from the aqueous media was much higher when the cells are induced by growth in HgCl_2 [150]. Tested environmental samples are inoculated in aqueous growth medium supplemented with known concentration of HgCl_2 and incubated for 24 hrs at 37°C. After centrifugation, the remaining concentration of mercury present in supernatant is measured using the Atomic Absorption Mercury Analyzer and compared to initial known concentration of Hg^{+2} to calculate its removal rate [83, 149].

Moreover, about 1ml samples are taken every 4 hrs and centrifuged. Then, remaining Hg concentration is measured in supernatants after processing. Hg^{+2} concentrations are calculated by measuring absorbance at OD600 using standard curve. Then, the initial and the final Hg^{+2} concentrations can be determined and reduction rates will be calculated [87].

8.2.3. Hg Volatilization Assay

The ability of bacteria to volatilize mercury from added mercuric chloride is a common mercury resistance mechanism by *mer* genes [150]. So, volatilization can be determined by using mercury vapor analyzer [75] or other atomic absorption spectrometry [78, 151], radioactive assay or using sensitive film containing Ag^+ as TEM photography film (X-ray film) which forms foggy areas due to reduction of the Ag^+ emulsion of the film by the mercury vapor [148, 151].

8.2.4. Mercuric Reductase (MR) Activity

MR activity measured depending on Hg^{+2} reduction mechanism by the enzyme in presence of NADPH. Hg^{2+} forms a complex with two cysteines of the enzyme active site. NADPH transfers a proton to FAD forming FADH. The resulting FADH then reduces Hg^{2+} into Hg, and oxidizes back into a FAD. After reduction, the mercury is then released from the enzyme as a volatile vapor [152].

The MR activity was determined by different methods

1. Fluorimetric measurements by oxidation of fluorescent NADPH to the non-fluorescent NADP^+ due to the reduction of Hg^{+2} to Hg at specific wave lengths [144].
2. MR assays by measuring Mercury-dependent NADPH oxidation using a UV-visible spectrophotometer [153, 154] as NADPH increases the activity of MR while NADP^+ has no effect on activity [154].
3. Determination of the remaining amount of NADPH by titration using phenazine methosulfate to produce visible formazan. The concentration of formed formazan is detected Spectrophotometric. As enzyme activity is related to the remaining amount of NADPH and the produced amount of formazan [155].

8.2.5. Biosorption Experiments

Analysis of control and inoculated samples for metal adsorption by inductively coupled plasma optical emission spectroscopy [156].

CONCLUSION

Industrialization has a great threat to the environmental mercury contamination already spread in the atmosphere,

soil and water systems. Mercury causes more damage when reached to humans and animals. So, mercury usage should be restricted more to reduce its pollution. *mer* genes enable bacteria to convert the toxic organic or inorganic mercury forms to less toxic forms helping in mercury bioremediation as the most environmental friendly, safe and effective remediation technique.

The main *mer* operon genes are the *merB* and *merA* genes both can help bacteria to detoxify both organic mercurial compounds and inorganic mercury. *merA*, is a mercuric ion reductase enzyme, catalyzes the reduction of Hg^{+2} into volatile Hg . *merB*, codes for organomercurial lyase catalyzes demethylation of organic mercury compounds. *merT*, *merP*, *merC*, *merE*, *merF*, and *merH* express different transporter proteins. *merG* helps in cellular effluxing of organomercurial compounds (Phenylmercury). *merR* encodes for a metalloregulatory protein, it is Hg^{+2} dependent transcriptional repressor-activator that can sense metal concentration and control the expression of other functional *mer* operon genes. Evolution and diversity in structure and organization of *mer* genes between different Gram-negative and Gram-positive strains affects their resistance ability as the transporter *merP* and *merT* genes were more common to occur in earliest evolved less complex operons while, other transporter genes as *merC*, *merF*, *merE*, *merH* and effluxing *merG* gene are commonly occurred in recently evolved, more complex operons [89].

Bacteria harboring the *mer* determinants can be genetically modified to code for resistance to other toxicants as toxic metals and be used during extreme environmental different conditions. *mer* determinants could be transferred into a new genetically engineered recipient suitable biological system (plant, bacteria or algae) to help in mercury remediation after removing undesired genes or adding genes to increase system harmony. Although *mer* system is highly understood mercury cycle in anaerobic conditions specially MeHg formation mechanism that prefers anoxic conditions is incompletely understood and requires more investigations as MeHg is the most bioaccumulative toxic form and understanding the anaerobic system will help in MeHg bioremediation.

CONFLICTS OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

CONSENT FOR PUBLICATION

Not applicable

ACKNOWLEDGEMENTS

The manuscript was drafted by Dr. Martha M. Naguib and revised and approved by Dr. Ahmed O. El-Gendy, Dr. Ahmed S. Khairalla respectively.

REFERENCES

- [1] Report IT. Heavy Metals ”— A meaningless term ? (IUPAC Technical Report). 2002; 74(5): 793-807.
- [2] Montuelle B, Latour X, Volat B, Gounot AM. Toxicity of heavy metals to bacteria in sediments. Bull Environ Contam Toxicol 1994; 53(5): 753-8.
[<http://dx.doi.org/10.1007/BF00196950>] [PMID: 7833613]
- [3] Nisa M, Coral Ü, Korkmaz H, Arikian B, Coral G. Plasmid mediated heavy metal resistances in Enterobacter spp. In: isolated from Sofulu landfill. in Adana, Turkey. Growth (Lakeland). 2005; 55: pp. (3)175-9.
- [4] Nies DH. Microbial heavy-metal resistance. Appl Microbiol Biotechnol 1999; 51(6): 730-50.
[<http://dx.doi.org/10.1007/s002530051457>] [PMID: 10422221]
- [5] Wedepohl KH. The composition of the continental crust. Geochim Cosmochim Acta 1995; 59(7): 1217-32.
[[http://dx.doi.org/10.1016/0016-7037\(95\)00038-2](http://dx.doi.org/10.1016/0016-7037(95)00038-2)]
- [6] Figueiredo NLL, Areias A, Mendes R, Canário J, Duarte A, Carvalho C. Mercury-resistant bacteria from salt marsh of Tagus Estuary: the influence of plants presence and mercury contamination levels. J Toxicol Environ Health A 2014; 77(14-16): 959-71.
[<http://dx.doi.org/10.1080/15287394.2014.911136>] [PMID: 25072727]
- [7] Brooks S, Moore C, Lew D, Lefer B, Huey G, Tanner D. Temperature and sunlight controls of mercury oxidation and deposition atop the Greenland ice sheet. Atmos Chem Phys 2011; 11(16): 8295-306.
[<http://dx.doi.org/10.5194/acp-11-8295-2011>]
- [8] Dash HR, Das S. Bioremediation of mercury and the importance of bacterial mer genes. Int Biodeterior Biodegradation 2012; 75: 207-13.
[<http://dx.doi.org/10.1016/j.ibiod.2012.07.023>]

- [9] Keating MH, Beauregard D, Benjey WG, *et al.* Mercury Study Report to Congress Volume II: An Inventory of Anthropogenic Mercury Emissions in the United States. United States Environmental Protection Agency 1998. EPA-452/R-1-181.
- [10] Dash HR, Das S. International biodeterioration & biodegradation bioremediation of mercury and the importance of bacterial mer genes. *Int Biodeterior Biodegradation* 2012; 75: 207-13.
[<http://dx.doi.org/10.1016/j.ibiod.2012.07.023>]
- [11] Barkay T, Miller SM, Summers AO. Bacterial mercury resistance from atoms to ecosystems. *FEMS Microbiol Rev* 2003; 27(2-3): 355-84.
[[http://dx.doi.org/10.1016/S0168-6445\(03\)00046-9](http://dx.doi.org/10.1016/S0168-6445(03)00046-9)] [PMID: 12829275]
- [12] Wang Q, Kim D, Dionysiou DD, Sorial Ga, Timberlake D. Sources and remediation for mercury contamination in aquatic systems - A literature review. *Environ Pollut* 2004; 131(2): 323-36.
[<http://dx.doi.org/10.1016/j.envpol.2004.01.010>] [PMID: 15234099]
- [13] Smith T, Pitts K, McGarvey JA, Summers AO. Bacterial oxidation of mercury metal vapor, Hg(0). *Appl Environ Microbiol* 1998; 64(4): 1328-32.
[PMID: 9546169]
- [14] Lindqvist O, Rodhe H. Atmospheric mercury—A review. *Tellus B Chem Phys Meteorol* 1985; 37(3): 136-59.
[<http://dx.doi.org/10.3402/tellusb.v37i3.15010>]
- [15] Keeler G, Glinsorn G, Pirrone N. Particulate mercury in the atmosphere: Its significance, transport, transformation and sources. *Water Air Soil Pollut* 1995; 80(1-4): 159-68.
[<http://dx.doi.org/10.1007/BF01189664>]
- [16] Munthe J, McElroy WJ. Some aqueous reactions of potential importance in the atmospheric chemistry of mercury. *Atmos Environ Part A* 1992; 26(4): 553-7.
[[http://dx.doi.org/10.1016/0960-1686\(92\)90168-K](http://dx.doi.org/10.1016/0960-1686(92)90168-K)]
- [17] Matthew J, Colomboa JH, John R. Reinfelder, Tamar Barkayb, Nathan Yeea. Anaerobic oxidation of Hg(0) and methylmercury formation by *Desulfovibrio desulfuricans* ND132. *Geochim Cosmochim Acta* 2013; 112: 166-77.
[<http://dx.doi.org/10.1016/j.gca.2013.03.001>]
- [18] Allard B, Arsenie I. Abiotic reduction of mercury by humic substances in aquatic system — An important process for the mercury cycle. *Water Air Soil Pollut* 1991; 56(1): 457-64.
[<http://dx.doi.org/10.1007/BF00342291>]
- [19] Colombo MJ, Ha J, Reinfelder JR, Barkay T, Yee N. Oxidation of Hg(0) to Hg(II) by diverse anaerobic bacteria. *Chem Geol* 2014; 363: 334-40.
[<http://dx.doi.org/10.1016/j.chemgeo.2013.11.020>]
- [20] Zheng W, Liang L, Gu B. Mercury reduction and oxidation by reduced natural organic matter in anoxic environments. *Environ Sci Technol* 2012; 46(1): 292-9.
[<http://dx.doi.org/10.1021/es203402p>] [PMID: 22107154]
- [21] Gu B, Bian Y, Miller CL, Dong W, Jiang X, Liang L. Mercury reduction and complexation by natural organic matter in anoxic environments. *Proc Natl Acad Sci USA* 2011; 108(4): 1479-83.
[<http://dx.doi.org/10.1073/pnas.1008747108>] [PMID: 21220311]
- [22] Hamlett NV, Landale EC, Davis BH, Summers AO. Roles of the Tn21 merT, merP, and merC gene products in mercury resistance and mercury binding. *J Bacteriol* 1992; 174(20): 6377-85.
[<http://dx.doi.org/10.1128/jb.174.20.6377-6385.1992>] [PMID: 1328156]
- [23] Erie L. Mechanistic steps in the photoreduction of mercury in natural waters. 1994; 9697(1993).
- [24] Skogerboe RK, Wilson SA. Reduction of ionic species by fulvic acid. *Anal Chem* 1981; 53(2): 228-32.
[<http://dx.doi.org/10.1021/ac00225a023>]
- [25] 2013.
- [26] Podar M, Gilmour CC, Brandt CC, *et al.* Global prevalence and distribution of genes and microorganisms involved in mercury methylation. *Sci Adv* 2015; 1(9): e1500675.
[<http://dx.doi.org/10.1126/sciadv.1500675>] [PMID: 26601305]
- [27] Parks JM, Johs A, Podar M, *et al.* The genetic basis for bacterial mercury methylation. *Science* 2013; 339(6125): 1332-5.
[<http://dx.doi.org/10.1126/science.1230667>] [PMID: 23393089]
- [28] Sedimentt AE. Principal methylators 1985; 50(2): 498-502.
- [29] Wood JM, Kennedy FS, Rosen CG. Synthesis of methyl-mercury compounds by extracts of a methanogenic bacterium. *Nature* 1968; 220(5163): 173-4.
[<http://dx.doi.org/10.1038/220173a0>] [PMID: 5693442]
- [30] Choi SC, Chase T, Bartha R. Metabolic pathways leading to mercury methylation in *Desulfovibrio desulfuricans* LS. *Appl Environ Microbiol* 1994; 60(11): 4072-7.
[PMID: 16349435]

- [31] Hamelin S, Amyot M, Barkay T, Wang Y, Planas D. Methanogens: Principal methylators of mercury in lake periphyton. *Environ Sci Technol* 2011; 45(18): 7693-700. [http://dx.doi.org/10.1021/es2010072] [PMID: 21875053]
- [32] Kerin EJ, Gilmour CC, Roden E, Suzuki MT, Coates JD, Mason RP. Mercury methylation by dissimilatory iron-reducing bacteria. *Appl Environ Microbiol* 2006; 72(12): 7919-21. [http://dx.doi.org/10.1128/AEM.01602-06] [PMID: 17056699]
- [33] Hu H, Lin H, Zheng W, *et al.* Oxidation and methylation of dissolved elemental mercury by anaerobic bacteria. *Nat Geosci* 2013; 6(9): 751-4. [http://dx.doi.org/10.1038/ngeo1894]
- [34] Li Y, Cai Y. Progress in the study of mercury methylation and demethylation in aquatic environments. *Chin Sci Bull* 2013; 58(2): 177-85. [http://dx.doi.org/10.1007/s11434-012-5416-4]
- [35] Cerrati G, Bernhard M, Weber JH. Model reactions for abiotic mercury (II) methylation: Kinetics of methylation of mercury (II) by mono□, di□, and tri□methyltin in seawater. *Appl Organomet Chem* 1992; 6(7): 587-95. [http://dx.doi.org/10.1002/aoc.590060705]
- [36] Li Y, Yin Y, Liu G, *et al.* Estimation of the major source and sink of methylmercury in the Florida Everglades. *Environ Sci Technol* 2012; 46(11): 5885-93. [http://dx.doi.org/10.1021/es204410x] [PMID: 22536798]
- [37] Begley TP, Waits AE, Walsh CT. Bacterial organomercurial Lyase: Over production, Isolation, and characterization? 1986(1982); 7186-92.
- [38] Oremland RS, Culbertson CW, Winfrey MR. Methylmercury decomposition in sediments and bacterial cultures: Involvement of methanogens and sulfate reducers in oxidative demethylation. *Appl Environ Microbiol* 1991; 57(1): 130-7. [PMID: 16348388]
- [39] Schaefer JK, Letowski J, Barkay T. mer-mediated resistance and volatilization of Hg (II) under anaerobic conditions. *Geomicrobiol J* 2002; 19(1): 87-102. [http://dx.doi.org/10.1080/014904502317246192]
- [40] Summers AO, Silver S. Mercury resistance in a plasmid-bearing strain of *Escherichia coli*. *J Bacteriol* 1972; 112(3): 1228-36. [PMID: 4565536]
- [41] Lu X, Liu Y, Johns A, *et al.* Anaerobic mercury methylation and demethylation by geobacter bemidjensis Bem. *Environ Sci Technol* 2016; 50(8): 4366-73. [http://dx.doi.org/10.1021/acs.est.6b00401] [PMID: 27019098]
- [42] Tjerngren I, Karlsson T, Björn E, Skjellberg U. Potential Hg methylation and MeHg demethylation rates related to the nutrient status of different boreal wetlands. *Biogeochem* 2012; 108(1): 335-50. [http://dx.doi.org/10.1007/s10533-011-9603-1]
- [43] Suda I, Suda M, Hirayama K. Degradation of methyl and ethyl mercury by singlet oxygen generated from sea water exposed to sunlight or ultraviolet light. *Arch Toxicol* 1993; 67(5): 365-8. [http://dx.doi.org/10.1007/BF01973709] [PMID: 8368946]
- [44] Seller P, Kelly CA, Rudd JWM, MacHutchon AR. Photodegradation of methylmercury in lakes. *Nature* 1996; 380(6576): 694-7. [http://dx.doi.org/10.1038/380694a0]
- [45] Naftz DL, Cederberg JR, Krabbenhoft DP, Beisner KR, Whitehead J, Gardberg J. Diurnal trends in methylmercury concentration in a wetland adjacent to Great Salt Lake, Utah, USA. *Chem Geol* 2011; 283(1-2): 78-86. [http://dx.doi.org/10.1016/j.chemgeo.2011.02.005]
- [46] Nascimento AM, Chartone-Souza E. Operon mer: Bacterial resistance to mercury and potential for bioremediation of contaminated environments. *Genet Mol Res* 2003; 2(1): 92-101. [PMID: 12917805]
- [47] Braune B, Chételat J, Amyot M, *et al.* Mercury in the marine environment of the Canadian Arctic: review of recent findings. *Sci Total Environ* 2015; 509-510: 67-90. [http://dx.doi.org/10.1016/j.scitotenv.2014.05.133] [PMID: 24953756]
- [48] Dixit R, Wasiullah EY, Malaviya D, *et al.* Bioremediation of heavy metals from soil and aquatic environment: An overview of principles and criteria of fundamental processes. *Sustainability* 2015; 7(2): 2189-212. [http://dx.doi.org/10.3390/su7022189]
- [49] Deng X, Wang P. Isolation of marine bacteria highly resistant to mercury and their bioaccumulation process. *Bioresour Technol* 2012; 121: 342-7. [http://dx.doi.org/10.1016/j.biortech.2012.07.017] [PMID: 22864169]
- [50] Rizwan M, Singh M, Mitra CK, Morve RK. Ecofriendly application of nanomaterials: Nanobioremediation. *J Nanoparticles* 2014; 2014: 7. [http://dx.doi.org/10.1155/2014/431787]
- [51] McCarthy D, Edwards GC, Gustin MS, Care A, Miller MB, Sunna A. An innovative approach to bioremediation of mercury contaminated soils from industrial mining operations. *Chemosphere* 2017; 184: 694-9. [http://dx.doi.org/10.1016/j.chemosphere.2017.06.051] [PMID: 28633064]

- [52] Elekes CC, Busuioc G. The mycoremediation of metals polluted soils using wild growing species of mushrooms. Jul 22; 2010; pp. In: In Proceedings of the 7th WSEAS international conference on Latest trends on Eng Educ, Corfu Island Greece 2010; 36-9.
- [53] Wang J, Feng X, Anderson CWN, Xing Y, Shang L. Remediation of mercury contaminated sites - A review. *J Hazard Mater* 2012; 221-222: 1-18.
[<http://dx.doi.org/10.1016/j.jhazmat.2012.04.035>] [PMID: 22579459]
- [54] Ruiz ON, Daniell H. Genetic engineering to enhance mercury phytoremediation. *Curr Opin Biotechnol* 2009; 20(2): 213-9.
[<http://dx.doi.org/10.1016/j.copbio.2009.02.010>] [PMID: 19328673]
- [55] Sone Y, Nakamura R, Pan-Hou H, Sato MH, Itoh T, Kiyono M. Increase methylmercury accumulation in *Arabidopsis thaliana* expressing bacterial broad-spectrum mercury transporter MerE. *AMB Express* 2013; 3(1): 52.
[<http://dx.doi.org/10.1186/2191-0855-3-52>] [PMID: 24004544]
- [56] Rugh CL, Wilde HD, Stack NM, Thompson DM, Summers AO, Meagher RB. Mercuric ion reduction and resistance in transgenic *Arabidopsis thaliana* plants expressing a modified bacterial merA gene. *Proc Natl Acad Sci USA* 1996; 93(8): 3182-7.
[<http://dx.doi.org/10.1073/pnas.93.8.3182>] [PMID: 8622910]
- [57] Bizily SP, Rugh CL, Summers AO, Meagher RB. Phytoremediation of methylmercury pollution: merB expression in *Arabidopsis thaliana* confers resistance to organomercurials. *Proc Natl Acad Sci USA* 1999; 96(12): 6808-13.
[<http://dx.doi.org/10.1073/pnas.96.12.6808>] [PMID: 10359794]
- [58] Rugh CL, Senecoff JF, Meagher RB, Merkle SA. Development of transgenic yellow poplar for mercury phytoremediation. *Nat Biotechnol* 1998; 16(10): 925-8.
[<http://dx.doi.org/10.1038/nbt1098-925>] [PMID: 9788347]
- [59] He YK, Sun JG, Feng XZ, Czakó M, Márton L. Differential mercury volatilization by tobacco organs expressing a modified bacterial merA gene. *Cell Res* 2001; 11(3): 231-6.
[<http://dx.doi.org/10.1038/sj.cr.7290091>] [PMID: 11642409]
- [60] Ruiz ON, Hussein HS, Terry N, Daniell H. Phytoremediation of organomercurial compounds via chloroplast genetic engineering. *Plant Physiol* 2003; 132(3): 1344-52.
[<http://dx.doi.org/10.1104/pp.103.020958>] [PMID: 12857816]
- [61] Yang H, Nairn J, Ozias-Akins P, Ozias-Akins P. Transformation of peanut using a modified bacterial mercuric ion reductase gene driven by an actin promoter from *Arabidopsis thaliana*. *J Plant Physiol* 2003; 160(8): 945-52.
[<http://dx.doi.org/10.1078/0176-1617-01087>] [PMID: 12964870]
- [62] Czakó M, Feng X, He Y, Liang D, Márton L. Transgenic *Spartina alterniflora* for phytoremediation. *Environ Geochem Health* 2006; 28(1-2): 103-10.
[<http://dx.doi.org/10.1007/s10653-005-9019-8>] [PMID: 16528587]
- [63] Heaton AC, Rugh CL, Kim T, Wang NJ, Meagher RB. Toward detoxifying mercury-polluted aquatic sediments with rice genetically engineered for mercury resistance. *Environ Toxicol Chem* 2003; 22(12): 2940-7.
[<http://dx.doi.org/10.1897/02-442>] [PMID: 14713034]
- [64] Che D, Meagher RB, Heaton AC, Lima A, Rugh CL, Merkle SA. Expression of mercuric ion reductase in Eastern cottonwood (*Populus deltoides*) confers mercuric ion reduction and resistance. *Plant Biotechnol J* 2003; 1(4): 311-9.
[<http://dx.doi.org/10.1046/j.1467-7652.2003.00031.x>] [PMID: 17163907]
- [65] Lyyra S, Meagher RB, Kim T, et al. Coupling two mercury resistance genes in Eastern cottonwood enhances the processing of organomercury. *Plant Biotechnol J* 2007; 5(2): 254-62.
[<http://dx.doi.org/10.1111/j.1467-7652.2006.00236.x>] [PMID: 17309680]
- [66] Nagata T, Ishikawa C, Kiyono M, Pan-Hou H. Accumulation of mercury in transgenic tobacco expressing bacterial polyphosphate. *Biol Pharm Bull* 2006; 29(12): 2350-3.
[<http://dx.doi.org/10.1248/bpb.29.2350>] [PMID: 17142961]
- [67] Sinha A, Khare SK. Mercury bioaccumulation and simultaneous nanoparticle synthesis by *Enterobacter* sp. cells. *Bioresour Technol* 2011; 102(5): 4281-4.
[<http://dx.doi.org/10.1016/j.biortech.2010.12.040>] [PMID: 21216593]
- [68] Kretsinger RH, Uversky VN, Permyakov EA. *Encyclopedia of Metalloproteins*. Springer 2013.
[<http://dx.doi.org/10.1007/978-1-4614-1533-6>]
- [69] Baldi F, Pepi M, Filippelli M. Methylmercury resistance in *Desulfovibrio desulfuricans* strains in relation to methylmercury degradation. *Appl Environ Microbiol* 1993; 59(8): 2479-85.
[PMID: 16349013]
- [70] Bhakta V, Balkrishna M, Thakuri C. Bacterial mer operon-mediated detoxification of mercurial compounds: A short review. *Arch Microbiol* 2011; 837-44.
- [71] Pan-Hou HS, Nishimoto M, Imura N. Possible role of membrane proteins in mercury resistance of *Enterobacter aerogenes*. *Arch Microbiol* 1981; 130(2): 93-5.
[<http://dx.doi.org/10.1007/BF00411057>] [PMID: 6459062]

- [72] Dash HR, Das S. Bioremediation of mercury and the importance of bacterial mer genes. *Int Biodeterior Biodegradation* 2012; 75: 207-13. [<http://dx.doi.org/10.1016/j.ibiod.2012.07.023>]
- [73] Sugio T, Komoda T, Okazaki Y, Takeda Y, Nakamura S, Takeuchi F. Volatilization of metal mercury from Organomercurials by highly mercury-resistant *Acidithiobacillus ferrooxidans* MON-1. *Biosci Biotechnol Biochem* 2010; 74(5): 1007-12. [<http://dx.doi.org/10.1271/bbb.90888>] [PMID: 20460735]
- [74] Zeng XX, Tagn JX, Jiang P, Liu H W, Dai Z-mM, Liu X-dD. Isolation, characterization and extraction of mer gene of Hg²⁺ resisting strain D2. *Trans Nonferrous Met Soc China* 2010; 20(50621063): 507-12. [English Edition]. [[http://dx.doi.org/10.1016/S1003-6326\(09\)60170-9](http://dx.doi.org/10.1016/S1003-6326(09)60170-9)]
- [75] Schelert J, Dixit V, Hoang V, Simbahan J, Drozda M, Blum P. Occurrence and characterization of mercury resistance in the hyperthermophilic archaeon *Sulfolobus solfataricus* by use of gene disruption. *J Bacteriol* 2004; 186(2): 427-37. [<http://dx.doi.org/10.1128/JB.186.2.427-437.2004>] [PMID: 14702312]
- [76] Liebert CA, Hall RM, Summers AO. Transposon Tn21, flagship of the floating genome. *Microbiol Mol Biol Rev* 1999; 63(3): 507-22. [PMID: 10477306]
- [77] Wireman J, Liebert CA, Smith T, *et al.* Association of mercury resistance with antibiotic resistance in the gram-negative fecal bacteria of primates. *Association of Mercury Resistance with Antibiotic Resistance in the Gram-Negative Fecal Bacteria of Primates* 1997; 63(11)
- [78] Pepi M, Gaggi C, Bernardini E, *et al.* Mercury-resistant bacterial strains *Pseudomonas* and *Psychrobacter* spp. isolated from sediments of Orbetello Lagoon (Italy) and their possible use in bioremediation processes. *Int Biodeterior Biodegradation* 2011; 65(1): 85-91. [<http://dx.doi.org/10.1016/j.ibiod.2010.09.006>]
- [79] Figueiredo NLL, Canário J, Duarte A, Serralheiro ML, Carvalho C. Isolation and characterization of mercury-resistant bacteria from sediments of Tagus Estuary (Portugal): Implications for environmental and human health risk assessment. *J Toxicol Environ Health A* 2014; 77(1-3): 155-68. [<http://dx.doi.org/10.1080/15287394.2014.867204>] [PMID: 24555656]
- [80] D SR. Prevalence of Mercury-Resistant and Antibiotic-Resistant Bacteria found in Dental Amalgam. 2014; 3(4): 1-4.
- [81] Summers AO, Wireman J, Vimy MJ, *et al.* Mercury released from dental “silver” fillings provokes an increase in mercury- and antibiotic-resistant bacteria in oral and intestinal floras of primates. *Antimicrob Agents Chemother* 1993; 37(4): 825-34. [<http://dx.doi.org/10.1128/AAC.37.4.825>] [PMID: 8280208]
- [82] Ball MM, Carrero P, Castro D, Yarzabal LA. Mercury resistance in bacterial strains isolated from tailing ponds in a gold mining area near El Callao (Bolívar State, Venezuela). *Curr Microbiol* 2007; 54(2): 149-54. [<http://dx.doi.org/10.1007/s00284-006-0347-4>] [PMID: 17200804]
- [83] François F, Lombard C, Guigner JM, *et al.* Isolation and characterization of environmental bacteria capable of extracellular biosorption of mercury. *Appl Environ Microbiol* 2012; 78(4): 1097-106. [<http://dx.doi.org/10.1128/AEM.06522-11>] [PMID: 22156431]
- [84] De J, Ramaiah N. Characterization of marine bacteria highly resistant to mercury exhibiting multiple resistances to toxic chemicals. *Ecol Indic* 2007; 7(3): 511-20. [<http://dx.doi.org/10.1016/j.ecolind.2006.05.002>]
- [85] Bogdanova ES, Bass IA, Minakhin LS, *et al.* Horizontal spread of mer operons among gram-positive bacteria in natural environments. *Microbiology* 1998; 144(Pt 3): 609-20. [<http://dx.doi.org/10.1099/00221287-144-3-609>] [PMID: 9534232]
- [86] Dash HR, Sahu M, Mallick B, Das S. Functional efficiency of MerA protein among diverse mercury resistant bacteria for efficient use in bioremediation of inorganic mercury. *Biochimie* 2017; 142: 207-15. [<http://dx.doi.org/10.1016/j.biochi.2017.09.016>] [PMID: 28966143]
- [87] Yu Z, Li J, Li Y, *et al.* A mer operon confers mercury reduction in a *Staphylococcus epidermidis* strain isolated from Lanzhou reach of the Yellow River. *Int Biodeterior Biodegradation* 1981; 90: 57-63. [<http://dx.doi.org/10.1016/j.ibiod.2014.02.002>]
- [88] Naik MM, Dubey S. Lead-and mercury-resistant marine bacteria and their application in lead and mercury bioremediation *Marine Pollution and Microbial Remediation*. Springer 2017; pp. 29-40.
- [89] Boyd ES, Barkay T. The mercury resistance operon: from an origin in a geothermal environment to an efficient detoxification machine. *Front Microbiol* 2012; 3: 349. [<http://dx.doi.org/10.3389/fmicb.2012.00349>] [PMID: 23087676]
- [90] Sone Y, Nakamura R, Pan-hou H, Itoh T, Kiyono M. MerP in resistance to mercurials and the transport of mercurials in *Escherichia coli*. 2013; 36:1835-41.
- [91] Jan AT, Azam M, Ali A, Haq QMR. Molecular characterization of mercury resistant bacteria inhabiting polluted water bodies of different geographical locations in India. *Curr Microbiol* 2012; 65(1): 14-21. [<http://dx.doi.org/10.1007/s00284-012-0118-3>] [PMID: 22488489]
- [92] Wilson JR, Leang C, Morby AP, Hobman JL, Brown NL. MerF is a mercury transport protein: different structures but a common mechanism for mercuric ion transporters? *FEBS Lett* 2000; 472(1): 78-82. [[http://dx.doi.org/10.1016/S0014-5793\(00\)01430-7](http://dx.doi.org/10.1016/S0014-5793(00)01430-7)] [PMID: 10781809]

- [93] Kiyono M, Oka Y, Sone Y, *et al.* Bacterial heavy metal transporter MerC increases mercury accumulation in *Arabidopsis thaliana*. *Biochem Eng J* 2013; 71: 19-24.
[<http://dx.doi.org/10.1016/j.bej.2012.11.007>]
- [94] Schuë M, Dover LG, Besra GS, Parkhill J, Brown NL. Sequence and analysis of a plasmid-encoded mercury resistance operon from *Mycobacterium marinum* identifies MerH, a new mercuric ion transporter. *J Bacteriol* 2009; 191(1): 439-44.
[<http://dx.doi.org/10.1128/JB.01063-08>] [PMID: 18931130]
- [95] Schelert J, Rudrappa D, Johnson T, Blum P. Role of MerH in mercury resistance in the archaeon *Sulfolobus solfataricus*. *Microbiology* 2013; 159(Pt 6): 1198-208.
[<http://dx.doi.org/10.1099/mic.0.065854-0>] [PMID: 23619003]
- [96] Kiyono M, Pan-Hou H. The merG gene product is involved in phenylmercury resistance in *Pseudomonas* strain K-62. *J Bacteriol* 1999; 181(3): 726-30.
[PMID: 9922233]
- [97] Osborn AM, Bruce KD, Strike P, Ritchie DA. Distribution, diversity and evolution of the bacterial mercury resistance (mer) operon. *FEMS Microbiol Rev* 1997; 19(4): 239-62.
[<http://dx.doi.org/10.1111/j.1574-6976.1997.tb00300.x>] [PMID: 9167257]
- [98] Das S, Dash HR, Chakraborty J. Genetic basis and importance of metal resistant genes in bacteria for bioremediation of contaminated environments with toxic metal pollutants. *Appl Microbiol Biotechnol* 2016; 100(7): 2967-84.
[<http://dx.doi.org/10.1007/s00253-016-7364-4>] [PMID: 26860944]
- [99] Schottel JL. The mercuric and organomercurial detoxifying enzymes from a plasmid-bearing strain of *Escherichia coli*. *J Biol Chem* 1978; 253(12): 4341-9.
[PMID: 350872]
- [100] Fermentation BK, Inage K, Chiba T, Metallic J, G S. Metallic in mercury-resistant pseudomonas has not and the electron described is containing group, and the enzyme for the decomposition mercuric chloride on bag, and subjected to a column (2.5 x 25 cm) of DEAE. 1971; 36(12): 217-6.
- [101] Essa AMM. The effect of a continuous mercury stress on mercury reducing community of some characterized bacterial strains. *Afr J Microbiol Res* 2012; 6(18): 4006-12.
- [102] Pan-Hou H. [Application of mercury-resistant genes in bioremediation of mercurials in environments]. *Yakugaku Zasshi* 2010; 130(9): 1143-56.
[<http://dx.doi.org/10.1248/yakushi.130.1143>] [PMID: 20823672]
- [103] Parks JM, Guo H, Momany C, *et al.* Mechanism of Hg-C protonolysis in the organomercurial lyase MerB. *J Am Chem Soc* 2009; 131(37): 13278-85.
[<http://dx.doi.org/10.1021/ja9016123>] [PMID: 19719173]
- [104] Mathema VB, Krishna B, Thakuri C, Sillanpää M, Amatya R. Study of mercury (II) chloride tolerant bacterial isolates from Baghmati River with estimation of plasmid size and growth variation for the high mercury (II) resistant *Enterobacter* spp. *Culture* 2011; 72-.
- [105] Sone Y, Mochizuki Y, Koizawa K, *et al.* Mercurial-resistance determinants in *Pseudomonas* strain K-62 plasmid pMR68. *AMB Express* 2013; 3(1): 41.
[<http://dx.doi.org/10.1186/2191-0855-3-41>] [PMID: 23890172]
- [106] Guo H-B, Johs A, Parks JM, *et al.* Structure and conformational dynamics of the metalloregulator MerR upon binding of Hg(II). *J Mol Biol* 2010; 398(4): 555-68.
[<http://dx.doi.org/10.1016/j.jmb.2010.03.020>] [PMID: 20303978]
- [107] Park S-j, Wireman JOY, Summers A. Genetic Analysis of the Tn21 operator-promoter 1992; 174(7): 2160-71.
- [108] Chang C-C, Lin L-Y, Zou X-W, Huang C-C, Chan N-L. Structural basis of the mercury(II)-mediated conformational switching of the dual-function transcriptional regulator MerR. *Nucleic Acids Res* 2015; 43(15): 7612-23.
[<http://dx.doi.org/10.1093/nar/gkv681>] [PMID: 26150423]
- [109] Haberstroh L, Silver S. 1984.
- [110] Lee IW, Gambill BD, Summers AO. Translation of merD in Tn21. *J Bacteriol* 1989; 171(4): 2222-5.
[<http://dx.doi.org/10.1128/jb.171.4.2222-2225.1989>] [PMID: 2539363]
- [111] Wireman J, Liebert CA, Smith T, Summers AO. Association of mercury resistance with antibiotic resistance in the gram-negative fecal bacteria of primates. *Appl Environ Microbiol* 1997; 63(11): 4494-503.
[PMID: 9361435]
- [112] Zinder ND, Lederberg J. Genetic exchange in *Salmonella*. *J Bacteriol* 1952; 64(5): 679-99.
[PMID: 12999698]
- [113] Lederberg J. Gene recombination in the bacterium *Escherichia Coli*. *Nature* 1947; 673-84.
- [114] Frost LS, Leplae R, Summers AO, Toussaint A. Mobile genetic elements: the agents of open source evolution. *Nat Rev Microbiol* 2005; 3(9): 722-32.
[<http://dx.doi.org/10.1038/nrmicro1235>] [PMID: 16138100]

- [115] Dash HR, Das S. Diversity, community structure, and bioremediation potential of mercury-resistant marine bacteria of estuarine and coastal environments of Odisha, India. *Environ. Sci. Pollut Res.* 2015.
- [116] Rankin DJ, Rocha EPC, Brown SP. What traits are carried on mobile genetic elements, and why? *Heredity (Edinb)* 2011; 106(1): 1-10. [<http://dx.doi.org/10.1038/hdy.2010.24>] [PMID: 20332804]
- [117] Dimitriu T, Misevic D, Lindner AB, Taddei F. Mobile genetic elements are involved in bacterial sociality. *Mob Genet Elements* 2015; 5(1): 7-11. [<http://dx.doi.org/10.1080/2159256X.2015.1006110>] [PMID: 26435881]
- [118] Martinez JL. The role of natural environments in the evolution of resistance traits in pathogenic bacteria. *Proc Biol Sci* 2009; 276(1667): 2521-30. [<http://dx.doi.org/10.1098/rspb.2009.0320>] [PMID: 19364732]
- [119] Martínez JL. Antibiotics and antibiotic resistance genes in natural environments. *Science* 2008; 321(5887): 365-7. [<http://dx.doi.org/10.1126/science.1159483>] [PMID: 18635792]
- [120] Bennett PM. Genome plasticity: insertion sequence elements, transposons and integrons, and DNA rearrangement. *Methods Mol Biol* 2004; 266: 71-113. [PMID: 15148416]
- [121] Mindlin S, Kholodii G, Gorlenko Z, *et al.* Mercury resistance transposons of gram-negative environmental bacteria and their classification. *Res Microbiol* 2001; 152(9): 811-22. [[http://dx.doi.org/10.1016/S0923-2508\(01\)01265-7](http://dx.doi.org/10.1016/S0923-2508(01)01265-7)] [PMID: 11763242]
- [122] Bogdanova E, Minakhin L, Bass I, Volodin A, Hobman JL, Nikiforov V. Class II broad-spectrum mercury resistance transposons in Gram-positive bacteria from natural environments. *Res Microbiol* 2001; 152(5): 503-14. [[http://dx.doi.org/10.1016/S0923-2508\(01\)01224-4](http://dx.doi.org/10.1016/S0923-2508(01)01224-4)] [PMID: 11446519]
- [123] Matsui K, Yoshinami S, Narita M, *et al.* Mercury resistance transposons in Bacilli strains from different geographical regions. *FEMS Microbiol Lett* 2016; 363(5): fnw013. [<http://dx.doi.org/10.1093/femsle/fnw013>] [PMID: 26802071]
- [124] Liebert CA, Hall RM, Summers AO. Transposon Tn 21. Flagship of the Floating Genome 1999; 63(3): 507-22.
- [125] Lal D, Lal R. Evolution of mercuric reductase (merA) gene: A case of horizontal gene transfer. *Mikrobiologija* 2010; 79(4): 524-31. [PMID: 21058506]
- [126] Hall JPI, Harrison E, Lilley AK, Paterson S, Spiers AJ, Brockhurst MA. Environmentally co-occurring mercury resistance plasmids are genetically and phenotypically diverse and confer variable context-dependent fitness effects. *Environ Microbiol* 2015; 17(12): 5008-22. [<http://dx.doi.org/10.1111/1462-2920.12901>] [PMID: 25969927]
- [127] Huang CC, Narita M, Yamagata T, Endo G. Identification of three merB genes and characterization of a broad-spectrum mercury resistance module encoded by a class II transposon of *Bacillus megaterium* strain MB1. *Gene* 1999; 239(2): 361-6. [[http://dx.doi.org/10.1016/S0378-1119\(99\)00388-1](http://dx.doi.org/10.1016/S0378-1119(99)00388-1)] [PMID: 10548738]
- [128] Laddaga RA, Chu L, Misra TK, Silver S. Nucleotide sequence and expression of the mercurial-resistance operon from *Staphylococcus aureus* plasmid pI258. *Proc Natl Acad Sci USA* 1987; 84(15): 5106-10. [<http://dx.doi.org/10.1073/pnas.84.15.5106>] [PMID: 3037534]
- [129] Petrovski S, Blackmore DW, Jackson KL, Stanisich VA. Mercury(II)-resistance transposons Tn502 and Tn512, from *Pseudomonas* clinical strains, are structurally different members of the Tn5053 family. *Plasmid* 2011; 65(1): 58-64. [<http://dx.doi.org/10.1016/j.plasmid.2010.08.003>] [PMID: 20800080]
- [130] Osborn AM, Bruce KD, Ritchie DA, Strike P. The mercury resistance operon of the IncJ plasmid pMERPH exhibits structural and regulatory divergence from other Gram-negative mer operons. *Microbiology* 1996; 142(Pt 2): 337-45. [<http://dx.doi.org/10.1099/13500872-142-2-337>] [PMID: 8932707]
- [131] Narita M, Chiba K, Nishizawa H, *et al.* Diversity of mercury resistance determinants among Bacillus strains isolated from sediment of Minamata Bay. *FEMS Microbiol Lett* 2003; 223(1): 73-82. [[http://dx.doi.org/10.1016/S0378-1097\(03\)00325-2](http://dx.doi.org/10.1016/S0378-1097(03)00325-2)] [PMID: 12799003]
- [132] Wang Y, Moore M, Levinson HS, Silver S, Walsh C, Mahler I. Nucleotide sequence of a chromosomal mercury resistance determinant from a *Bacillus* sp. with broad-spectrum mercury resistance. *J Bacteriol* 1989; 171(1): 83-92. [<http://dx.doi.org/10.1128/jb.171.1.83-92.1989>] [PMID: 2536669]
- [133] Griffin HG, Foster TJ, Silver S, Misra TK. Cloning and DNA sequence of the mercuric- and organomercurial-resistance determinants of plasmid pDU1358. *Proc Natl Acad Sci USA* 1987; 84(10): 3112-6. [<http://dx.doi.org/10.1073/pnas.84.10.3112>] [PMID: 3033633]
- [134] Reniero D, Galli E, Barbieri P. Cloning and comparison of mercury- and organomercurial-resistance determinants from a *Pseudomonas stutzeri* plasmid. *Gene* 1995; 166(1): 77-82. [[http://dx.doi.org/10.1016/0378-1119\(95\)00546-4](http://dx.doi.org/10.1016/0378-1119(95)00546-4)] [PMID: 8529897]
- [135] Kholodii GY, Mindlin SZ, Bass IA, Yurieva OV, Minakhina SV, Nikiforov VG. Four genes, two ends, and a res region are involved in transposition of Tn5053: A paradigm for a novel family of transposons carrying either a mer operon or an integron. *Mol Microbiol* 1995; 17(6): 1189-200.

- [http://dx.doi.org/10.1111/j.1365-2958.1995.mmi_17061189.x] [PMID: 8594337]
- [136] Christopher M, Paul O, Hamadi B. Association of metal tolerance with multidrug resistance among Environmental Bacteria from wetlands of Lake Victoria Basin 2014.
- [137] Zeyauallah M, Islam B, Ali a. Isolation, identification and PCR amplification of merA gene from highly mercury polluted Yamuna river. *Afr J Biotechnol* 2010; 9(24): 3510-4.
- [138] Pal C, Bengtsson-Palme J, Kristiansson E, Larsson DGJ. Co-occurrence of resistance genes to antibiotics, biocides and metals reveals novel insights into their co-selection potential. *BMC Genomics* 2015; 16: 964. [http://dx.doi.org/10.1186/s12864-015-2153-5] [PMID: 26576951]
- [139] Trajanovska S, Britz ML, Bhave M. Detection of heavy metal ion resistance genes in gram-positive and gram-negative bacteria isolated from a lead-contaminated site. *Biodegradation* 1997; 8(2): 113-24. [http://dx.doi.org/10.1023/A:1008212614677] [PMID: 9342884]
- [140] Diels L, Mergey M. DNA probe-mediated detection of resistant bacteria from soils highly polluted by heavy metals. *Appl Environ Microbiol* 1990; 56(5): 1485-91. [PMID: 16348196]
- [141] Bruce KD, Hiorns WD, Hobman JL, Osborn AM, Strike P, Ritchie DA. Amplification of DNA from native populations of soil bacteria by using the polymerase chain reaction. *Appl Environ Microbiol* 1992; 58(10): 3413-6. [PMID: 1444376]
- [142] Georgios M, Egki T. Phenotypic and Molecular Methods for the Detection of Antibiotic Resistance Mechanisms in Gram Negative Nosocomial Pathogens. Intech 2014. [http://dx.doi.org/10.5772/57582]
- [143] Barkay T, Fouts DL, Olson BH. Preparation of a DNA gene probe for detection of mercury resistance genes in gram-negative bacterial communities. *Appl Environ Microbiol* 1985; 49(3): 686-92. [PMID: 3994373]
- [144] Baldi F, Gallo M, Marchetto D, Faleri C, Maida I, Fani R. Manila clams from Hg polluted sediments of Marano and Grado lagoons (Italy) harbor detoxifying Hg resistant bacteria in soft tissues. *Environ Res* 2013; 125: 188-96. [http://dx.doi.org/10.1016/j.envres.2012.11.008] [PMID: 23398778]
- [145] Unc A, Zurek L, Peterson G, Narayanan S, Springthorpe SV, Sattar SA. Microarray assessment of virulence, antibiotic, and heavy metal resistance in an agricultural watershed creek. *J Environ Qual* 2012; 41(2): 534-43. [http://dx.doi.org/10.2134/jeq2011.0172] [PMID: 22370416]
- [146] Osborn AM, Bruce KD, Strike P, Ritchie DA. Polymerase Chain Reaction-Restriction Fragment Length Polymorphism Analysis Shows Divergence among mer Determinants from Gram-Negative Soil Bacteria Indistinguishable by DNA-DNA Hybridization 1993.
- [147] N. Mirzazai FK, Kargar M. Isolation and identification of mercury resistant bacteria from Kor river, Iran. 2008; pp. 935-9.
- [148] Taylor P, Figueiredo NLL, Areias A, Mendes R, Canário J, Duarte A. Mercury-Resistant Bacteria From Salt Marsh of Tagus Estuary: The Influence of Plants Presence and Mercury Contamination Levels. *J Toxicol Environ Health* 2014; •••: 37-41.
- [149] Cao D-j, Tian Z-f. Isolation and identification of a mercury resistant strain. *Environ Prot Eng* 2012; 38(4)
- [150] Summers AO, Lewis E. Volatilization of mercuric chloride by mercury-resistant plasmid-bearing strains of *Escherichia coli*, *Staphylococcus aureus*, and *Pseudomonas aeruginosa*. *J Bacteriol* 1973; 113(2): 1070-2. [PMID: 4632313]
- [151] Nakamura K, Nakahara H. Simplified X-ray film method for detection of bacterial volatilization of mercury chloride by *Escherichia coli*. *Appl Environ Microbiol* 1988; 54(11): 2871-3. [PMID: 3063210]
- [152] Lian P, Guo H-b, Riccardi D, Dong A, Parks JM, Xu Q. X-ray Structure of a Hg²⁺ Complex of Mercuric Reductase (MerA) and Quantum Mechanical/Molecular Mechanical Study of Hg²⁺ Transfer between the C-Terminal and Buried Catalytic Site Cysteine Pairs. 2014.
- [153] Freedman Z, Zhu C, Barkay T. Mercury resistance and mercuric reductase activities and expression among chemotrophic thermophilic Aquificae. *Appl Environ Microbiol* 2012; 78(18): 6568-75. [http://dx.doi.org/10.1128/AEM.01060-12] [PMID: 22773655]
- [154] Sandström A, Lindskog S. Activation of mercuric reductase by the substrate NADPH. *Eur J Biochem* 1987; 164(1): 243-9. [http://dx.doi.org/10.1111/j.1432-1033.1987.tb11017.x] [PMID: 3104042]
- [155] Ogunseitan OA. Protein method for investigating mercuric reductase gene expression in aquatic environments. *Appl Environ Microbiol* 1998; 64(2): 695-702. [PMID: 9464410]
- [156] Umrانيا VV. Bioremediation of toxic heavy metals using acidothermophilic autotrophes. *Bioresour Technol* 2006; 97(10): 1237-42. [http://dx.doi.org/10.1016/j.biortech.2005.04.048] [PMID: 16324838]
- [157] Bizily SP, Rugh CL, Summers AO, Meagher RB. Phytoremediation of methylmercury pollution: merB expression in *Arabidopsis thaliana* confers resistance to organomercurials. *Proc Natl Acad Sci USA* 1999; 96(12): 6808-13. [http://dx.doi.org/10.1073/pnas.96.12.6808] [PMID: 10359794]

- [158] Bizily SP, Rugh CL, Meagher RB. Phytodetoxification of hazardous organomercurials by genetically engineered plants. *Nat Biotechnol* 2000; 18(2): 213-7.
[<http://dx.doi.org/10.1038/72678>] [PMID: 10657131]
- [159] Hsieh J-L, Chen C-Y, Chiu M-H, *et al.* Expressing a bacterial mercuric ion binding protein in plant for phytoremediation of heavy metals. *J Hazard Mater* 2009; 161(2-3): 920-5.
[<http://dx.doi.org/10.1016/j.jhazmat.2008.04.079>] [PMID: 18538925]
- [160] Sasaki Y, Hayakawa T, Inoue C, Miyazaki A, Silver S, Kusano T. Generation of mercury-hyperaccumulating plants through transgenic expression of the bacterial mercury membrane transport protein MerC. *Transgenic Res* 2006; 15(5): 615-25.
[<http://dx.doi.org/10.1007/s11248-006-9008-4>] [PMID: 16830224]
- [161] Schelert J, Drozda M, Dixit V, Dillman A, Blum P. Regulation of mercury resistance in the crenarchaeote *Sulfolobus solfataricus*. *J Bacteriol* 2006; 188(20): 7141-50.
[<http://dx.doi.org/10.1128/JB.00558-06>] [PMID: 17015653]

© 2018 Naguib *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: (<https://creativecommons.org/licenses/by/4.0/legalcode>). This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.