

# Rapid *In Vitro* Regeneration and Genetic Fidelity Assessment of Regenerated Plants in *Ayapana Triplinervis* (Vahl) R.M. King & H. Robinson: An Ethnomedicinal and Ornamental Herb



Thumadath Palayullaparambil Ajeesh Krishna<sup>1</sup>, Neenthamadathil Mohandas Krishnakumar<sup>1,2,\*</sup>, Theivanayagam Maharajan<sup>1</sup> and Stanislaus Antony Ceasar<sup>1,2,\*</sup>

<sup>1</sup>Division of Plant Molecular Biology and Biotechnology, Department of Biosciences, Rajagiri College of Social Sciences (Autonomous), Kochi, Kerala 683 104, India

<sup>2</sup>Division of Phytochemistry and Drug-Design, Department of Biosciences, Rajagiri College of Social Sciences (Autonomous), Kochi, Kerala 683 104, India

## Abstract:

**Background::** *Ayapana triplinervis* is a popular ethnomedicinal and ornamental plant species. Due to its high medicinal importance, *A. triplinervis* was recently documented in the French Pharmacopeia.

**Objective::** Rapid and efficient tissue culture protocol development is crucial for the high production and biotechnological applications of this plant.

**Methods::** In this study, an efficient tissue culture protocol was developed for plant regeneration using nodal explants of *A. triplinervis*. The nodal explants were treated in Murashige and Skoog's (MS) medium supplemented with various individual concentrations of cytokinins (BAP and KIN) and auxins (IAA and IBA). The nodal explant was regenerated in three different steps: (1) initial shoot induction, (2) shoot multiplication and elongation, and (3) rooting.

**Results::** The results revealed that all individual concentrations (10, 20, 30, or 40 mg/L) of BAP or KIN responded to induce shoot initiation. The highest shoot multiplication and elongation were achieved in the MS medium with 20 mg/L BAP and 20 mg/L KIN. The regenerated plantlets produced better roots on MS medium containing 1.0 mg/L of each IAA or IBA. The well-established rooted plantlets were maintained in the culture room and greenhouse for better acclimatization and achieved a 100% survival rate. We analyzed the genetic fidelity of *in vitro* regenerated plants using random amplified polymorphic DNA (RAPD) markers. No genetic polymorphisms were observed *in vitro* plants compared to the mother plants.

**Conclusion::** This efficient protocol could benefit future biotechnological applications like mass multiplication, genetic transformation and gene editing for improving the bioactive molecules in *A. triplinervis*.

**Keywords:** *Ayapana triplinervis*, Genetic fidelity, Nodal explant, Phytohormones, Tissue culture, Plants.

© 2024 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

\*Address correspondence to these authors at the Division of Plant Molecular Biology and Biotechnology, Department of Biosciences, Rajagiri College of Social Sciences (Autonomous), Kochi-683104, Kerala, India; E-mails: [antony\\_sm2003@yahoo.co.in](mailto:antony_sm2003@yahoo.co.in) and [krishnakumarmohandas@gmail.com](mailto:krishnakumarmohandas@gmail.com)

Cite as: Krishna T, Krishnakumar N, Maharajan T, Ceasar S. Rapid *In Vitro* Regeneration and Genetic Fidelity Assessment of Regenerated Plants in *Ayapana Triplinervis* (Vahl) R.M. King & H. Robinson: An Ethnomedicinal and Ornamental Herb. Open Biotechnol J, 2024; 18: e18740707192866. <http://dx.doi.org/10.2174/0118740707192866240116112512>



CrossMark

Received: September 22, 2023

Revised: December 08, 2023

Accepted: December 27, 2023

Published: ?? ??, 2024



Send Orders for Reprints to [reprints@benthamscience.net](mailto:reprints@benthamscience.net)

## 1. INTRODUCTION

*Ayapana triplinervis* (Vahl) R. M. King & H. Robinson is a widely used folk medicinal plant in the healing processes, and it belongs to the family *Asteraceae*. The plant is used in mystical-religious rituals by the traditional communities inhabiting Asia, Africa, and South America [1]. It is an erect, perennial, semi-woody at basal region, ornamental herb growing up to a height of 1 m [2]. The leaves of the plant are widely used as decoctions, teas, or baths against viral infections and respiratory and gynecological diseases in Brazil, Bangladesh, Sri Lanka, West Indies, Mauritius, Peru, and some European countries [3]. It is commonly known as 'Ayapana' in India and 'Vishalyakarani' in Ayurvedic literature, and the plant is used in wounds, bleeding stomachs, and diarrhoea because of its capacity to control blood coagulation [3]. The plant is also a good cardiac stimulant, emetic, laxative, expectorant, alterative, and antiscorbutic and is used in cases of dyspepsia and agues [4].

The phytochemical investigations on *A. triplinervis* by the researchers revealed the presence of secondary metabolites, such as coumarins (ayapin, ayapanin, daphnetin etc.), essential oils (thymoquinone, thymohydroquinone dimethyl ether,  $\beta$ -caryophyllene,  $\beta$ -selinene, dimethyl ether thymoquinone and 2, 5-dimethoxy p-cymene), stigmasterol, and carotene [2, 5]. The pharmacological properties of *A. triplinervis* have been extensively studied, and the plant was found to possess various biological activities, such as hepatoprotective, antioxidant, antimicrobial, anti-nociceptive, anti-inflammatory, antimelanogenic, anxiolytic, antidepressant, antiulcer, anticancer, hypocholesteremic, gastroprotective, and antidiabetic effects [6-16]. A recent study revealed that thymohydroquinone dimethyl ether extracted from the aerial parts of *A. triplinervis* inhibited Zika virus infection in human epithelial A549 cells [17]. *A. triplinervis* is one of the most important medicinal plants used in herbal drug development. Ayapin, or 6, 7-methylenedioxy coumarin is a phytoalexin used as an antimicrobial and antiviral agent, was isolated from this plant [18]. The plant contains a variety of coumarins with an anticoagulant activity that are the precursors for drugs, such as 'Warfarin' [19]. The nano-emulsions developed from the essential oils extracted from *A. triplinervis* are used as mosquito repellent [20, 21].

According to the World Health Organization (WHO), the current demand for medicinal plants is increasing annually, estimated at about 14 billion US dollars [22]. The pharmaceutical industries mainly depend on the wild population of medicinal plants to extract phyto compounds in herbal drug development [23, 24]. However, the commercial management of these phytoconstituents is impeded due to the inadequate supply of medicinal plants [25]. The high demand for medicinal plants motivated rural farmers to start the commercial cultivation of medicinal plants. Previous attempts have been made for the *in vitro* propagation of *A. triplinervis* (Synonyms: *Eupatorium triplinerve*) using different explants. The micropropagation protocol through axillary bud prolifera-

tion and *ex vitro* rooting was reported [26]. Usha and Karpagam [27] developed a regeneration protocol using nodal explants supplemented with 4.44  $\mu$ M 6-Benzylaminopurine (BAP) and produced 18 shoots with an average length of 6.4 cm. In another study, more shoots (5) with higher shoot length (2.85 cm) were obtained from nodal explants on MS medium fortified with 0.2 mg/L BAP and 0.02 mg/L Gibberellic acid 3 ( $GA_3$ ) after 30 days. In another protocol, MS medium fortified with 17.76  $\mu$ M/L BAP and 2.69  $\mu$ M/L of naphthalene acetic acid (NAA) produced 4.93 shoots [28]. These studies showed that the low concentration of phytohormones had poor responses to shoot initiation and multiplication. The optimum concentration of phytohormones could help for better shoot initiation and multiplication. The optimization of phytohormone concentration is essential for the development of an efficient plant regeneration protocol. In this study, the previously reported same plant species *A. triplinervis* was used to develop an *in vitro* regenerative protocol. The study describes a rapid regenerative protocol of *A. triplinervis* using nodal explants for a maximum of 35 shoot induction with the assessment of the genetic fidelity of regenerated plantlets using RAPD markers. This is the first study reporting maximum shoot induction in this plant species. This study may be useful for mass multiplication and biotechnological studies of *A. triplinervis*.

## 2. MATERIALS AND METHODS

### 2.1. Plant Material

*A. triplinervis* plants were collected from the Rajagiri College of Social Sciences (Autonomous) herbal garden, Kakkanad campus, Kochi, Kerala, India, and a herbarium was prepared for morphological study. Herbarium specimens were taxonomically identified and authenticated by a botanist. Previously reported plant *A. triplinervis* (Synonyms: *Eupatorium triplinerve*) was used for this present study [27]. The protocol described by Antony Ceasar *et al.* [29] was used to prepare nodal explants with minor modifications in the surface sterilization procedure. Nodal explants (approximately 2 cm, each containing two opposite nodes) were washed in sterile water for 15 min and soaked in 0.1% (w/v) Bavistin containing carbendazim (BASF, Mumbai, India) for 20 min. It was then washed three times using sterile distilled water. Surface sterilization was performed by immersing nodal explants in 70% (v/v) alcohol for 10 seconds and washing three times using autoclaved distilled water. After that, the explants were treated with 0.1% (w/v)  $HgCl_2$  for 1 min and washed four times using sterile distilled water. The nodal explants were air-dried on filter paper (Whatman No. 1, Code:1001-125, Sigma-Aldrich) in the laminar airflow chamber. The sterilized nodal explants were used for further studies.

### 2.2. Culture Medium and Conditions

The MS basal medium (Himedia, Code: PT099-25L) was used in the present study [30]. It contained basal salts and vitamins, 3% sucrose (w/v); finally, 0.1 M NaOH was

used to adjust the pH 5.6-5.8 of the medium before adding 0.8% agar (w/v) to it. The medium was autoclaved for 15 min at 121°C and 15 lbs. Different concentrations of plant growth regulators were added to the basal MS medium before adding agar. The standardization and shoot induction studies were carried out in test tubes. After that, shoot multiplication, elongation, and root induction tests were performed in tissue culture bottles (6 x 13 cm). All cultures were sub-cultured in the same medium unless otherwise mentioned at 15-day intervals. All cultures were maintained under controlled conditions. The temperature was maintained at  $25 \pm 2^\circ\text{C}$  and 16/8 light/dark photoperiod using fluorescent tube lights with the light intensity of  $50 \mu\text{mol}/\text{m}^2/\text{s}$  photosynthetic photon flux density (PPFD).

### 2.3. Shoot Multiplication and Elongation

For shoot induction, nodal explants were cultured on an MS medium supplemented with different plant growth regulators (PGRs), such as BAP and kinetin (KIN). The experiment was performed as part of a standardization procedure to choose the optimum concentration of individual cytokines for efficient induction of shoots. Briefly, different concentrations (10, 20, 30, or 40 mg/L) of BAP and KIN were individually added to the MS medium. Three nodal explants were inoculated in each tissue culture bottle, and three replicates were maintained for each concentration. The optimum individual cytokine concentration was determined based on the number of shoots produced after three weeks of incubation in the light. The percentage of explants that responded and the number of shoots produced by the responding explants were calculated after three weeks of incubation. The three-week-old shoot cultures obtained from various concentrations of shoot induction medium were subcultured in the same multiplication medium for further shoot multiplication and elongation. The MS medium devoid of PGR was considered a control. The mean number of shoots and their length were recorded after four weeks of incubation in the light. The responses of shoot induction were calculated using the following formula [44].

$$\text{Response of shoot induction (\%)} = \frac{\text{Number of explants responded}}{\text{Total number of explant tested}}$$

### 2.4. Root Induction

The elongated shoots from the shoot clumps were excised aseptically and transferred to the rooting medium

amended individually with 0.5, 1.0, 1.5, or 2.0 mg/L indole-3-acetic acid (IAA) or indolebutyric acid (IBA) for rooting. The total number of roots produced per shoot and the mean length of the roots were measured after four weeks of incubation in the light. The responses of root induction were calculated using the following formula [44].

$$\text{Response of root induction (\%)} = \frac{\text{Number of explants responded}}{\text{Total number of explant tested}}$$

### 2.5. Acclimatization and Hardening

The rooted *in vitro* plants were cleaned in running tap water before being transplanted into paper cups containing sterilized horticultural-grade perlite (Astra Chemical, Chennai, India) and fed with 1/10 dilution liquid MS medium. The paper cups containing *in vitro* plantlets were covered with a transparent polythene bag to maintain humidity. These plants were maintained in the culture room for two weeks under the atmospheric conditions of  $25 \pm 2^\circ\text{C}$  temperature; light 16 h photoperiod with the light intensity of  $50 \mu\text{mol}/\text{m}^2/\text{s}$  PPFD; relative humidity,  $85 \pm 5\%$ . The 1/10 diluted liquid medium was supplied once in three days. After two weeks, the plantlets were removed from the cup, transferred to the pots containing soil and vermicompost in 1:1 proportion (v/v), and fed with tap water. Again, these plants were kept in the greenhouse for three days. Finally, the acclimatized plants were transferred to the field outside the greenhouse. The regenerated plants were grown *ex vivo* for about three months, and their survival rate and morphological characteristics were observed.

### 2.6. Genetic Fidelity Analysis

Genetic fidelity of *in vitro* regenerated plantlets of *A. triplinervis* with that of *ex-vitro* mother plants was confirmed by genomic DNA (gDNA) PCR amplification using random amplified polymorphic DNA (RAPD) primers (Table 1). The gDNA was isolated from *in vitro* regenerated plants of *A. triplinervis* (five plantlets were randomly selected) from *ex-vitro* along with the mother plants. Briefly, gDNA was extracted from a young leaves tissue sample using HiPurA® Plant DNA isolation kit (CTAB method) according to the manufacturer (Himedia, HiGenoMB, Cat. No: MB502-50PR) instructions. The quantity of gDNA was measured using a NanoDrop spectrophotometer (ND-ONE-W, Thermo Fisher Scientific, USA), and the quality of gDNA was checked via a 0.8% (w/v) agarose gel electrophoresis (BIO-RAD, Mini-Sub® Cell GT, USA).

**Table 1. Details of RAPD primers used for screening the DNA banding pattern in the mother plant of *A. triplinervis* and *in vitro* regenerated plantlets. The details of the primer code, primer sequence, and annealing temperature are provided.**

Sl. No.	Primer Code	Primer Sequence (5' to 3')	Annealing Tm (°C)
1	OPE-04	GTGACATGCC	40°C
2	OPF-05	CCGAATTCCC	40°C
3	OPR-07	ACTGGCCTGA	40°C

(Table 1) contd....

Sl. No.	Primer Code	Primer Sequence (5' to 3')	Annealing Tm (°C)
4	OPG-07	GAACCTGCCGG	40°C
5	OPK-10	GTGCAACGTG	40°C
6	OPO-14	AGCATGGCTC	40°C
7	OPW-18	TTCAGGGCAC	40°C

The PCR was set with 25 µL reaction mixture containing 100 ng gDNA, 1X Hi-Chrom PCR master mix, *Taq* DNA polymerase [1U], dNTPs [200 µM], and MgCl<sub>2</sub> [1 mM]) (Himedia, HiGenoMB, Cat. No: MBT089-50R) and 400 nM RAPD primer. The PCR was performed using a Thermal Cycler (BIO-RAD-T100, USA) with an initial denaturation at 95°C for 5 min, followed by 35 cycles of 30 s denaturation at 95°C, 30 s annealing at 40°C and 1 min extension at 72°C, with a final extension at 72°C for 7 min. The amplicons were resolved in a 2% (w/v) agarose gel electrophoresis (BIO-RAD, Mini-Sub® Cell GT, USA) and stained with ethidium bromide (EtBr). The resolved agarose gels were visualized using the GelDoc Go Imaging system (BIO-RAD, USA).

For data analysis, the amplification pattern of *in vitro* regenerated plantlets was compared with their mother plant. The amplification products were categorized into four classes, namely: (1) strong band and easy to score (2) weaker band, but able to score; (3) very weak band and challenging to score; and (4) no bands. Classes 1 and 2 were considered positive amplification and taken for the analysis, whereas classes 3 and 4 were considered negative for amplification and excluded from the scoring.

### 2.7. Statistical Analysis

Each treatment consisted of three replicates, and all the experiments were repeated thrice. The data were presented as mean ± SD using SPSS version 17.0 SPSS (SPSS Inc., Chicago, IL, USA), and the values were


considered significant at  $P < 0.05$ .

## 3. RESULTS AND DISCUSSION


### 3.1. Effect of Cytokinins on Shoot Multiplication and Elongation

Cytokinins are indispensable for developing multiple shoots [31]. They are the critical components of plant tissue culture. This study evaluated nodal explants of *A. triplinervis* on MS medium with two types of cytokinins (BAP and KIN) at various individual concentrations (10, 20, 30, or 40 mg/L) for developing rapid shoot multiplication protocol. The individual concentration of both BAP and KIN induced the shoot proliferation from the nodal stem segments of *A. triplinervis* after three weeks of *in vitro* culture. While MS medium without cytokinins (control) also influenced the growth of nodal segment buds, they did not show any shoot multiplication (Fig. 1). The percentage (%) of shoot induction ranged from 60–90% and 60–100% on MS medium with BAP and KIN, respectively (Table 2). The highest percentage for shoot initiation, 90% and 100%, was observed on MS medium containing 20 mg/L BAP and 20 mg/L KIN, respectively. It was followed by the medium containing 30 mg/L BAP and KIN, which was 80% (Table 2). After the shoot initiation, sub-culturing of the shoot clumps on the respective medium enhanced shoot multiplication (Fig. 2). Eight-week passages on this culture medium resulted in better shoot multiplication and elongation.

**Fig. (1).** Effect of types of phytohormones on the shoot regeneration of *A. triplinervis*. (A) Control-MS basal; (B) MS basal containing 20 mg/L KIN and (C) MS basal containing 20 mg/L BAP.



**Fig. (2).** Effect of different concentrations of kinetin (KIN) on the shoot regeneration in *A. triplinervis*. (A) Control-MS basal; (B) MS basal containing 10 mg/L KIN; (C) MS basal containing 20 mg/L KIN; (D) MS basal containing 30 mg/L KIN; and (E) MS basal containing 40 mg/L KIN.



**Fig. (3).** Plant regeneration from nodal explant of *A. triplinervis*. (A) The mother plant of habitat; (B) Nodal bud explants were used for micro-propagation; (C and D) Shoot induction on MS medium containing 20 mg/L BAP; (E) Induction of shooting after six weeks under controlled conditions (F) Shoot elongation on MS medium containing 20 mg/L BAP; (G) Elongated shoots inoculated in the MS medium containing 1.0 mg/L IAA; (H) Fully developed plantlets after eight weeks incubation (MS medium + 20 mg/L BAP + 1.0 mg/L IAA); (I) Acclimatized plantlets in sterile perlite at lab conditions and (J) Well-established plantlets in the natural environment.

**Table 2. Effect of cytokinins on shoot proliferation in *A. triplinervis* after eight weeks' incubation.**

Growth Regulators (mg/L)		Response of Shoot Initiation (%)	Number of Shoots/Explants	Shoot Length (cm)
KIN	00	00	00.00 ± 0.00	00.00 ± 0.00
	10	70	17.00 ± 1.73 <sup>d</sup>	05.20 ± 0.52 <sup>b</sup>
	20	100	31.66 ± 2.51 <sup>a</sup>	06.06 ± 0.30 <sup>a,b</sup>
	30	80	27.33 ± 1.52 <sup>b</sup>	07.26 ± 0.41 <sup>a</sup>
	40	60	20.66 ± 1.15 <sup>c</sup>	05.50 ± 0.43 <sup>b</sup>
BAP	00	00	00.00 ± 0.00	00.00 ± 0.00
	10	60	18.66 ± 1.52 <sup>d</sup>	06.46 ± 0.41 <sup>a</sup>
	20	90	35.33 ± 2.08 <sup>a</sup>	06.50 ± 0.88 <sup>a</sup>
	30	80	29.33 ± 0.57 <sup>b</sup>	05.90 ± 0.15 <sup>a</sup>
	40	60	22.00 ± 2.64 <sup>c</sup>	04.60 ± 0.35 <sup>a</sup>

**Note:** Values carrying the same alphabet did not vary significantly (Tukey's HSD;  $P > 0.05$ ).

Among the two cytokinins (BAP and KIN), better shoot multiplication and elongation responses were observed in the medium containing BAP. In BAP supplementation, significantly higher mean shoot number (35.33) and length (6.50 cm) were observed at 20 mg/L BAP (Table 2 and Fig. 3). The medium containing 10 mg/L BAP showed the lowest shoot number (18.66) and length (4.60 cm). Similarly, MS medium supplemented with 20 mg/L KIN showed better shoot multiplication for KIN (Table 2). However, the highest shoot length (7.26 cm) was observed in the medium containing 30 mg/L KIN. Previously, Gandhi and Saravanan [28] analyzed the effect of the combined and individual cytokinins on the shoot initiation and multiplication in *A. triplinervis*. In particular, the highest shoot number (24) and length (6.50 cm) were observed at 5.15 mg/L BAP and 2.06 mg/L KIN, respectively. The combination of cytokinins (5.15 mg/L BAP + 0.62 mg/L KIN) showed a slight improvement in shoot multiplication (26.00) but did not have significant variations [28]. Therefore, our result showed better shoot multiplication at 20 mg/L BAP with 35 mean shoots per explant.

In contrast with our results, Gandhi and Saravanan [28] reported the highest shoot length (6.50 cm) at the lower concentration of BAP (2.06 mg/L) in *A. triplinervis* (Synonyms: *Eupatorium triplinerve*). Our results showed a similar shoot length (6.50 cm) was obtained at 20 mg/L BAP. Also, our results revealed that extremely low (10 mg/L) and high (40 mg/L) concentrations of BAP and KIN inhibited shoot number and length (Table 2). Previous studies indicated that supplementation of higher and lower concentrations of BAP/KIN inhibits shoot multiplication in *A. triplinervis* [19, 27, 28]. Borchetia et al. [32] observed that higher concentrations of BAP inhibited shoot growth (nodal explant) in tea plants. Supplementation of high concentrations of BAP/KIN could inhibit the adventitious meristem elongation; it may reduce the shoot number [32, 33]. Therefore, the optimum concentration is essential for better and rapid shoot multiplication. Based on our study, a medium containing 20 mg/L BAP helps rapidly multiply *A. triplinervis* than previously reported works.

### 3.2. Effect of Auxins on Root Induction

The root is the main organ of plants, and it has many roles and functions. Therefore, adventitious root initiation is the prerequisite for developing a successful tissue culture protocol. Auxins play a leading role in regulating the initial growth of adventitious roots [34]. In this study, the *in vitro* regenerated *A. triplinervis* shoots were separated and treated with various concentrations (0.5, 1.0, 1.5, or 2.0 mg/L) of auxins (IAA or IBA) individually for root induction. Root initiation was observed in all concentrations of both IAA and IBA after three weeks of *in vitro* culture. However, a longer (4-6 weeks) incubation period was needed for better rooting (number and length) of regenerated plantlets. MS medium supplemented with various concentrations of IAA or IBA produced 100% rooting responses (Table 3). The MS medium without auxins (control) also showed root formation in nodal segment buds. However, no root proliferation was shown in the stem (an immersed portion of the MS medium). The findings of the present work supported the results of Balakrishnan et al. [19], who also reported that *in vitro* raised shoots produced roots in the media used for shoot multiplication without adding auxins. This finding was missing in the previously reported tissue culture on *A. triplinervis* [28, 35]. Babaei et al. [36] revealed that roots of *Curculigo latifolia* explant were induced faster in MS medium (control) devoid of plant growth regulators than in various concentrations (0.25, 0.50, or 0.75 mg/L) of IBA. It may be due to sufficient endogenous auxins within the explants. These findings indicated that auxin might be unnecessary for *in vitro* root proliferation for *A. triplinervis*.

However, our results revealed that the concentrations of IAA and IBA significantly influenced root initiation. Among the two auxins (IAA and IBA), better initiation was observed with IAA. The MS medium containing 1.0 mg/L IAA showed higher root initiation with 13.66 mean roots per explant (Table 3 and Fig. 3). Similarly, best rooting was observed in MS medium supplemented with 1.0 mg/L IBA with 10.33 mean roots per explant having a mean root length of 4.76 cm (Table 3). In contrast with our results, Samyudurai et al. [35] reported that the half-strength MS medium with various concentrations of IAA and IBA (0.2,

0.4, 0.6, 0.8 or 1.0 mg/L) showed the lowest root number and length in *A. triplinervis*. But, Balakrishnan *et al.* [19] showed better root initiation (12.36) of *A. triplinervis* at MS basal with 1.0 mg/L IBA. The present study suggests that 1.0 mg/L IAA was the most suitable concentration for root induction of *A. triplinervis*.

### 3.3. Acclimatization and Hardening

The acclimatization of micro-propagated plants is the most precious work because regenerated plantlets are highly susceptible to environmental effects. In this study, the regenerated healthy plantlets were transplanted to an artificial soil mixture (perlite) in paper cups and maintained in a culture room with controlled growth conditions for 14 days for better acclimatization (Fig. 31). The well-established plantlets were successfully transferred to the *ex vitro* conditions; they showed good growth and stayed healthy. The regenerated plantlets achieved a 100% survival rate. Their shoots flourished, and leaves became bigger and greenish under field conditions. The plants produced by micropropagation were morphologically and physiologically similar to mother plants. Acclimating *in vitro*-derived plants to the external environment is crucial for *A. triplinervis* [27]. Gandhi and Saravanan [28] reported that the *in vitro* regenerated *A. triplinervis* plantlets showed a 90% survival rate. However, the present study showed a better acclimatization procedure for successfully surviving *in vitro* regenerated plantlets.

### 3.4. Genetic Fidelity Analysis of *In Vitro* Generated Plantlets

Genetic variations are common and severe *in vitro*-generated plantlets [37]. Therefore, evaluation of the

genetic variation among the *in vitro* regenerated and mother plants is essential. So, molecular markers are preferable for evaluating genetic fidelity analysis. This study analyzed the genetic fidelity of *in vitro* regenerated *A. triplinervis* plantlets and their mother plant using RAPD markers. Totally seven RAPD primers were used in the present study, and these primers generated 40 PCR amplification products ranging from 190 to 1600 bp size (Table 4). The primer OPW 18 amplified the maximum number of nine bands (Fig. 4c) and was followed by primer OPG 07 (Table 4). The OPE 04 and OPF 05 primers amplified the lowest number of bands (Fig. 4a and 4e). Our results showed that all primer banding patterns of the mother plant were similar to *in vitro* regenerated plants (Fig. 4). No RAPD polymorphism was observed in the *in vitro* regenerated plants. For example, the primer OPO 14 produced six amplicons ranging from 270-1150 bp (270, 360, 500, 610, 750, and 1150 bp) in the mother and their *in vitro* generated plantlets of *A. triplinervis*. These results revealed that the RAPD markers could be an easy way to check the quality of *in vitro* regenerated plants. Previously, the RAPD markers were used to analyze the genetic homogeneity of micropropagated plants. RAPD marker-based genetic fidelity has been successfully explored in many plants, such as *Teucrium polium* [38], *Citrus limon* [39], *Gloriosa superba* [40], *Bacopa monnieri* [29], *Flemingia macrophylla* [41], *Andrographis alata* [42], and *Tecoma stans* [43]. To the best of our knowledge, this is the first report on the genetic fidelity analysis of micropropagated plants of *A. triplinervis* using the RAPD markers. Our results show that RAPD markers can be used to gain rapid and precise information about genetic similarities *in vitro* generated plants.

**Table 3. Effect of auxins on root proliferation in *A. triplinervis* after six weeks' incubation. The details of auxin (KIN and BAP) concentration, root initiation (%), root numbers, and root length are provided.**

Growth Regulators (mg/L)		Response of Root Initiation (%)	Number of Root/Explant	Root Length (cm)
IAA	0	0	00.00 ± 0.00	00.00 ± 0.00
	0.5	100	08.00 ± 2.00 <sup>c</sup>	03.96 ± 0.25 <sup>a</sup>
	1	100	13.66 ± 1.15 <sup>a</sup>	04.36 ± 0.32 <sup>a</sup>
	1.5	100	12.33 ± 1.52 <sup>a,b</sup>	03.70 ± 0.25 <sup>a</sup>
	2	100	12.00 ± 1.73 <sup>b</sup>	03.40 ± 0.40 <sup>a</sup>
IBA	0	0	00.00 ± 0.00	0.00 ± 0.00
	0.5	100	08.66 ± 0.67 <sup>a,b</sup>	03.53 ± 0.30 <sup>a</sup>
	1	100	10.33 ± 2.30 <sup>a</sup>	04.76 ± 0.25 <sup>a</sup>
	1.5	100	07.00 ± 1.73 <sup>b</sup>	04.00 ± 0.20 <sup>a</sup>
	2	100	07.66 ± 0.57 <sup>b</sup>	03.70 ± 0.23 <sup>a</sup>


**Note:** Values carrying the same alphabet did not vary significantly (Tukey's HSD;  $P > 0.05$ ).

**Table 4. DNA fingerprinting of *in vitro* regenerated plantlets and their mother plants. The details on the number of scorable bands and approximate sizes of the amplified band size are presented.**

Primer Code	No. of the Amplified Band	Size range (bp)	Approximate Size (bp) of the Amplified Band
OPE 04	3	210-400	210, 350 and 400
OPF 05	3	190-360	190, 300, and 360

(Table 4) contd....

Primer Code	No. of the Amplified Band	Size range (bp)	Approximate Size (bp) of the Amplified Band
OPR 07	5	200-600	200, 310, 420, 480 and 600
OPG 07	8	320-780	320, 340, 440, 490, 550, 600, 700, and 780
OPK 10	6	270-1150	270, 360, 500, 610, 750 and 1150
OPO 14	6	330-1160	330, 470, 620, 710, 950, and 1160
OPW 18	9	310-1600	310, 350, 430, 480, 580, 650, 690, 950 and 1600
<b>Total</b>	<b>40</b>	<b>190-1600</b>	-



**Fig. (4).** Evaluation of the genetic fidelity of regenerated plantlets and their mother plant of *A. triplinervis* using RAPD marker. DNA fingerprinting patterns were generated with RAPD primers OPE 04 (a), OPF 07 (b), OPW 18 (c), OPO 14 (d), and OPF 05 (e). Lane 1, DNA markers (100/50 bp ladder); Lane 2, RAPD marker profile of a mother plant. Lanes 3-7, RAPD marker profiles of five separate regenerated plantlets.



## CONCLUSION

This study reports an efficient *in vitro* regeneration protocol for direct shoot regeneration from the nodal explants of *A. triplinervis*. The MS basal with 20 mg/L BAP showed the highest shoot induction and elongation responses. The better rooting was obtained at 1.0 mg/L IAA. This work reports the optimum concentration of phytohormones for rapid and efficient plant regeneration of *A. triplinervis*. The acclimatization of *in vitro*-generated plantlets achieved a 100% survival rate. We successfully determined the genetic fidelity of plantlets using RAPD molecular markers. All seven RAPD primers produced monomorphic bands, confirming the genetic homogeneity of the *in vitro*-regenerated plants. Moreover, this efficient protocol could benefit future transgenic and gene editing works for altering and improving the biosynthesis pathways for bioactive molecules from *A. triplinervis*.

## LIST OF BBREVIATIONS

BAP	=	6-Benzylaminopurine
GA <sub>3</sub>	=	Gibberellic acid 3
IAA	=	Indole-3-acetic Acid
IBA	=	Indolebutyric Acid
KIN	=	Kinetin
MS	=	Murashige and Skoog's
NAA	=	Naphthalene Acetic Acid
PGRs	=	Plant Growth Regulators
RAPD	=	Random Amplified Polymorphic DNA

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

## HUMAN AND ANIMAL RIGHTS

No animals/humans were used in this research.

## CONSENT FOR PUBLICATION

Not applicable.

## AVAILABILITY OF DATA AND MATERIALS

The data and supportive information are available within the article.

## FUNDING

This work was financially supported by Rajagiri College of Social Sciences (Autonomous), Kalamassery, under Seed Money for Faculty Minor Research to the second author (NMK).

## CONFLICT OF INTEREST

Dr. Antony Ceasar Stanislaus is on the Editorial Advisory Board of the journal The Open Biotechnology Journal.

## ACKNOWLEDGEMENTS

We sincerely thank Rajagiri College of Social Sciences

(Autonomous), Kochi, Kerala, for providing the research facilities and support.

## REFERENCES

- [1] Rodrigues ABL, Lopes de Matos J, Martins RL, et al. Ethnopharmacological use, secondary metabolites and biological activity of *ayapana triplinervis* (VAHL) R. M. King and H. Rob.: A systematic review. *Pharmacogn Rev* 2022; 16(32): 70-3. <http://dx.doi.org/10.5530/phrev.2022.16.10>
- [2] Gauvin-Bialecki A, Marodon C. Essential oil of *Ayapana triplinervis* from Reunion Island: A good natural source of thymohydroquinone dimethyl ether. *Biochem Syst Ecol* 2008; 36(11): 853-8. <http://dx.doi.org/10.1016/j.bse.2008.09.006>
- [3] Cheriyan BV, scarlet S, Priyadarshini J, Joshi S, Santhseelan M, Mohamed S. *Eupatorium triplinerve* (Vahl): An ethnobotanical review. *AJPSci* 2019; 9(3): 200-2. <http://dx.doi.org/10.5958/2231-5691.2019.00032.7>
- [4] Khare CP. Indian medicinal plants. Springer-Verlag Berlin 2007. <http://dx.doi.org/10.1007/978-0-387-70638-2>
- [5] Trang N, Wanner M, Phuong LV, Koomen G, Dung N. Thymoquinone from *Eupatorium ayapana*. *Planta Med* 1993; 59(1): 99-9. <http://dx.doi.org/10.1055/s-2006-959619> PMID: 17230346
- [6] Bose P, Gupta M, Mazumder UK, Kumar RS, Sivakumar T, Kumar RS. Hepatoprotective and antioxidant effects of *Eupatorium ayapana* against carbon tetrachloride induced hepatotoxicity in rats. *Iran J Pharmacol Ther* 2007; 6: 27-33.
- [7] Gupta M, Mazumder UK, Chaudhuri I, et al. Antimicrobial activity of *Eupatorium ayapana*. *Fitoterapia* 2002; 73(2): 168-70. [http://dx.doi.org/10.1016/S0367-326X\(02\)00007-2](http://dx.doi.org/10.1016/S0367-326X(02)00007-2) PMID: 11978434
- [8] Viet Huong DT, Giang PM, Trang VM. Coumarins and polar constituents from *Eupatorium triplinerve* and evaluation of their  $\alpha$ -glucosidase inhibitory activity. *J Chem* 2020; 2020: 1-8. <http://dx.doi.org/10.1155/2020/8945063>
- [9] Yadava RN, Saini VK. *In vitro* antimicrobial efficacy of the essential oil of *Eupatorium triplinerve* leaves. *Indian perfum* 1990; 34(1): 61-3.
- [10] Parimala K, Cheriyan BV, Viswanathan S. Antinociceptive and anti-inflammatory activity of petroleum-ether extract of *Eupatorium triplinerve* Vahl. *J Pharm Sci* 2012; 2(3): 12-8.
- [11] Arung ET, Kuspradini H, Kusuma IW, Shimizu K, Kondo R. Validation of *Eupatorium triplinerve* Vahl leaves, a skin care herb from East Kalimantan, using a melanin biosynthesis assay. *J Acupunct Meridian Stud* 2012; 5(2): 87-92. <http://dx.doi.org/10.1016/j.jams.2012.01.003> PMID: 22483187
- [12] Melo AS, Monteiro MC, da Silva JB, et al. Antinociceptive, neurobehavioral and antioxidant effects of *Eupatorium triplinerve* Vahl on rats. *J Ethnopharmacol* 2013; 147(2): 293-301. <http://dx.doi.org/10.1016/j.jep.2013.03.002> PMID: 23524186
- [13] Krishnan M, Jayaraj RL, Megala J, Elangovan N. Antioxidant mediated antiulcer effect of *Eupatorium triplinerve* Vahl against acetic acid induced ulcerative colitis in mice. *Biomed Aging Pathol* 2014; 4(2): 153-60.
- [14] Maity P, Choudhury SM, Bepari M. Antimitotic, apoptotic and antineoplastic potential of leaf extract of *Eupatorium ayapana*. *Int J Phytomed* 2015; 7(1): 69-77.
- [15] Selvamangai G, Bhaskar A. GC-MS analysis of phytochemicals in the methanolic extract of *Eupatorium triplinerve*. *Asian Pac J Trop Biomed* 2012; 2(3): S1329-32. [http://dx.doi.org/10.1016/S2221-1691\(12\)60410-9](http://dx.doi.org/10.1016/S2221-1691(12)60410-9)
- [16] Maity R, Mondal P, Giri MK, Ghosh C, Mallick C. Gastroprotective effect of hydromethanolic extract of *Ayapana triplinervis* leaves on indomethacin-induced gastric ulcer in male Wistar rats. *J Food Biochem* 2021; 45(8): e13859. <http://dx.doi.org/10.1111/jfbc.13859> PMID: 34258791
- [17] Haddad JG, Picard M, Bénard S, et al. *Ayapana triplinervis*

- essential oil and its main component thymohydroquinone dimethyl ether inhibit zika virus at doses devoid of toxicity in zebrafish. *Molecules* 2019; 24(19): 3447.  
<http://dx.doi.org/10.3390/molecules24193447> PMID: 31547527
- [18] Venugopala KN, Rashmi V, Odhav B. Review on natural coumarin lead compounds for their pharmacological activity. *BioMed Res Int* 2013; 2013: 1-14.  
<http://dx.doi.org/10.1155/2013/963248> PMID: 23586066
- [19] Balakrishnan A, Aswindas TP, Sebastian DP, George S. *In vitro* axillary bud multiplication of an important medicinal plant-*Ayapana triplinervis* Vahl. *Int J Botany Stud* 2021; 6(4): 329-32.
- [20] Lobato Rodrigues AB, Martins RL, Rabelo ÉM, et al. Development of nano-emulsions based on *Ayapana triplinervis* essential oil for the control of *Aedes aegypti* larvae. *PLoS One* 2021; 16(7): e0254225.  
<http://dx.doi.org/10.1371/journal.pone.0254225> PMID: 34242328
- [21] Rodrigues ABL, Martins RL, Rabelo ÉM, et al. *In silico* and *in vivo* study of adulticidal activity from *Ayapana triplinervis* essential oils nano-emulsion against *Aedes aegypti*. *Arab J Chem* 2022; 15(9): 104033.  
<http://dx.doi.org/10.1016/j.arabjc.2022.104033>
- [22] Singh A. A note on variation of active principles in Indian medicinal plants and TIM formulations. *Ethnobot leafl* 2008; 2008(1)
- [23] Adarsh Krishna TP, Ajeesh Krishna TP, Sanyo Raj VN, et al. Evaluation of phytochemical constituents and proximate contents of the ethanolic leaf extract of *Tetrastigma leucostaphylum* (Dennst.) Alstone (Vitaceae) found in Western Ghats of Kerala, India. *Res J Pharm Sci* 2013; 2(10): 1-6.
- [24] Ajeesh Krishna TP, Adarsh Krishna TP, Kumuthakallavalli R, et al. Physico-chemical evaluation and biochemical quantification of crude drug powder (stem) of *Chassalia curviflora* (Wall. Ex Kurz.) Thwaites; a folk medicinal plant. *J Pharmacogn Phytochem* 2014; 3: 121-4.
- [25] Tippani R, Vemunoori AK, Yarra R, Nanna RS, Abbagani S, Thammidala C. Adventitious shoot regeneration from immature zygotic embryos of Indian Kino tree (*Pterocarpus marsupium* Roxb.) and genetic integrity analysis of *in vitro* derived plants using ISSR markers. *Hortic Environ Biotechnol* 2013; 54(6): 531-7.  
<http://dx.doi.org/10.1007/s13580-013-0161-4>
- [26] Martin KP. Rapid axillary bud proliferation and *ex vitro* rooting of *Eupatorium triplinerve*. *Biol Plant* 2003; 46(4): 589-91.  
<http://dx.doi.org/10.1023/B:BIOP.0000041067.23890.58>
- [27] Usha S, Karpagam S. Plant regeneration from nodal explants culture through organogenesis in *Eupatorium triplinerve* Vahl. *IOSR J Pharm Biol Sci* 2017; 12: 28-32.  
<http://dx.doi.org/10.9790/3008-1204032832>
- [28] Gandhi S. *In vitro* regeneration of *Eupatorium triplinerve* Vahl- an important medicinal herb through axillary bud culture. *Int J Pharm Biol Sci* 2018; 8: 563-8.
- [29] Antony Ceasar S, Lenin Maxwell S, Bhargav Prasad K, Karthigan M, Ignacimuthu S. Highly efficient shoot regeneration of *Bacopa monnieri* (L.) using a two-stage culture procedure and assessment of genetic integrity of micropropagated plants by RAPD. *Acta Physiol Plant* 2010; 32(3): 443-52.  
<http://dx.doi.org/10.1007/s11738-009-0419-8>
- [30] Murashige T, Skoog F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol Plant* 1962; 15(3): 473-97.  
<http://dx.doi.org/10.1111/j.1399-3054.1962.tb08052.x>
- [31] Schaller GE, Street IH, Kieber JJ. Cytokinin and the cell cycle. *Curr Opin Plant Biol* 2014; 21: 7-15.  
<http://dx.doi.org/10.1016/j.pbi.2014.05.015> PMID: 24994531
- [32] Borchetia S, Das SC, Handique PJ, Das S. High multiplication frequency and genetic stability for commercialization of the three varieties of micropropagated tea plants (*Camellia* spp.). *Sci Hortic* 2009; 120(4): 544-50.  
<http://dx.doi.org/10.1016/j.scienta.2008.12.007>
- [33] Deo B, Keshari B, Pradhan B. *In vitro* propagation of popular banana cultivar (Musa spp. Cv. Patakपुरa). *Bangladesh J Agric Res* 2020; 44(4): 641-8.  
<http://dx.doi.org/10.3329/bjar.v44i4.45699>
- [34] Pacurar DI, Perrone I, Bellini C. Auxin is a central player in the hormone cross-talks that control adventitious rooting. *Physiol Plant* 2014; 151(1): 83-96.  
<http://dx.doi.org/10.1111/ppl.12171> PMID: 24547793
- [35] Samydrurai P, Sarvesan R, Eganathan P. *In vitro* multiple shoot induction of *Eupatorium triplinerve* Vahl through direct regeneration from nodal segments. *Int J Appl Bioresearch* 2012; 6: 19-22.
- [36] Babaei N, Psyquay ANA, Saleh G, Lee Abdullah T. An efficient *in vitro* plantlet regeneration from shoot tip cultures of *Curculigo latifolia*, a medicinal plant. *ScientificWorldJournal* 2014; 2014: 1-9.  
<http://dx.doi.org/10.1155/2014/275028> PMID: 24723799
- [37] Chirumamilla P, Gopu C, Jogam P, Taduri S. Highly efficient rapid micropropagation and assessment of genetic fidelity of regenerants by ISSR and SCOT markers of *Solanum khasianum* Clarke. *Plant Cell Tissue Organ Cult* 2021; 144(2): 397-407.  
<http://dx.doi.org/10.1007/s11240-020-01964-6>
- [38] Rad FA, Jafari M, Khezrinejad N, Miandoab MP. An efficient plant regeneration system via direct organogenesis with *in vitro* flavonoid accumulation and analysis of genetic fidelity among regenerants of *Teucrium polium* L. *Hortic Environ Biotechnol* 2014; 55(6): 568-77.  
<http://dx.doi.org/10.1007/s13580-014-0611-7>
- [39] Goswami K, Sharma R, Singh PK, Singh G. Micropropagation of seedless lemon (*Citrus limon* L. cv. Kaghzi Kalan) and assessment of genetic fidelity of micropropagated plants using RAPD markers. *Physiol Mol Biol Plants* 2013; 19(1): 137-45.  
<http://dx.doi.org/10.1007/s12298-012-0148-0> PMID: 24381445
- [40] Yadav K, Aggarwal A, Singh N. Evaluation of genetic fidelity among micropropagated plants of *Gloriosa superba* L. using DNA-based markers — a potential medicinal plant. *Fitoterapia* 2013; 89: 265-70.  
<http://dx.doi.org/10.1016/j.fitote.2013.06.009> PMID: 23811099
- [41] Sirikonda A, Jogam P, Ellendula R, Kudikala H, Mood K, Allini VR. *In vitro* micropropagation and genetic fidelity assessment in *Flemingia macrophylla* (Willd.) Merr: An ethnomedicinal plant. *Vegetos* 2020; 33(2): 286-95.  
<http://dx.doi.org/10.1007/s42535-020-00106-9>
- [42] Kadapatti SS, Murthy HN. Rapid plant regeneration, analysis of genetic fidelity, and neoandrographolide content of micropropagated plants of *Andrographis alata* (Vahl) Nees. *J Genet Eng Biotechnol* 2021; 19(1): 20.  
<http://dx.doi.org/10.1186/s43141-021-00122-5> PMID: 33496903
- [43] Hussain SA, Anis M, Alatar AA. Efficient *in vitro* regeneration system for *Tecoma stans* L., using shoot tip and assessment of genetic fidelity among regenerants. *Proc Natl Acad Sci, India, Sect B Biol Sci* 2020; 90(1): 171-8.  
<http://dx.doi.org/10.1007/s40011-019-01093-x>
- [44] Ray BP, Hassan L, Nasiruddin KM. *In vitro* regeneration of brinjal (*Solanum melongena* L.). *Bangladesh J Agric Res* 1970; 36(3): 397-406.  
<http://dx.doi.org/10.3329/bjar.v36i3.9268>